

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
3 January 2002 (03.01.2002)

PCT

(10) International Publication Number  
**WO 02/00856 A2**

- (51) International Patent Classification<sup>7</sup>: **C12N 9/00** **M.**; Ketelbuiserstraat 99, B-8810 Lichtervelde (BE).  
**GEYSENS, Steven, C., J.**; Steenbeekstraaje 8, B-9770  
Kruishoutem (BE).
- (21) International Application Number: PCT/IB01/01514
- (22) International Filing Date: 29 June 2001 (29.06.2001) (74) Agent: **PUST, Detlev, A.**; Hoffmann . Eitle, Arabellas-  
trasse 4, 81925 München (DE).
- (25) Filing Language: English (81) Designated States (national): AU, CA, JP, KR, MX, NO.
- (26) Publication Language: English (84) Designated States (regional): European patent (AT, BE,  
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,  
NL, PT, SE, TR).
- (30) Priority Data:  
60/215,676 30 June 2000 (30.06.2000) US  
Published:  
— without international search report and to be republished  
upon receipt of that report
- (71) Applicant: **FLANDERS INTERUNIVERSITY INSTI-  
TUTE FOR BIOTECHNOLOGY (VIB)** [BE/BE]; Ri-  
jvisschestraat 120, B-9052 Zwijnaarde (BE).  
*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*
- (72) Inventors: **CONTRERAS, Roland**; Molenstraat 53,  
B-9820 Merelbeke (BE). **CALLEWAERT, Nico, L.**,



**WO 02/00856 A2**

(54) Title: PROTEIN GLYCOSYLATION MODIFICATION IN *PICHLIA PASTORIS*

(57) Abstract: The present invention provides genetically engineered strains of *Pichia* capable of producing proteins with reduced glycosylation. In particular, the genetically engineered strains of the present invention are capable of expressing either or both of an  $\alpha$ -1,2-mannosidase and glucosidase II. The genetically engineered strains of the present invention can be further modified such that the OCH1 gene is disrupted. Methods of producing glycoproteins with reduced glycosylation using such genetically engineered strains of *Pichia* are also provided.

### Protein Glycosylation Modification in *Pichia Pastoris*

#### **Field of the Invention**

The present invention relates to methods and vectors useful for genetically  
5 modifying the glycosylation process in methylotrophic yeast strains for the purpose of  
producing glycoproteins with reduced glycosylation. The present invention further  
relates to methylotrophic yeast strains generated using the present methods and  
vectors, as well as glycoproteins produced from such genetically modified strains.

#### **10 Background of the Invention**

The methylotrophic yeasts including *Pichia pastoris* have been widely used  
for production of recombinant proteins of commercial or medical importance.  
However, production and medical applications of some therapeutic glycoproteins can  
be hampered by the differences in the protein-linked carbohydrate biosynthesis  
15 between these yeasts and the target organism such as a mammalian subject.

Protein N-glycosylation originates in the endoplasmic reticulum (ER),  
where an N-linked oligosaccharide ( $\text{Glc}_3\text{Man}_9\text{GlcNAc}_2$ ) assembled on dolichol (a lipid  
carrier intermediate) is transferred to the appropriate Asn of a nascent protein. This is  
an event common to all eukaryotic N-linked glycoproteins. The three glucose residues  
20 and one specific  $\alpha$ -1,2-linked mannose residue are removed by specific glucosidases  
and an  $\alpha$ -1,2-mannosidase in the ER, resulting in the core oligosaccharide structure,  
 $\text{Man}_8\text{GlcNAc}_2$ . The protein with this core sugar structure is transported to the Golgi  
apparatus where the sugar moiety undergoes various modifications. There are  
significant differences in the modifications of the sugar chain in the Golgi apparatus  
25 between yeast and higher eukaryotes.

In mammalian cells, the modification of the sugar chain proceeds via 3  
different pathways depending on the protein moiety to which it is added. That is, (1)  
the core sugar chain does not change; (2) the core sugar chain is changed by adding  
the N-acetylglucosamine-1-phosphate moiety ( $\text{GlcNAc-1-P}$ ) in UDP-N-acetyl

glucosamine (UDP-GlcNAc) to the 6-position of mannose in the core sugar chain, followed by removing the GlcNAc moiety to form an acidic sugar chain in the glycoprotein; or (3) the core sugar chain is first converted into  $\text{Man}_5\text{GlcNAc}_2$  by removing 3 mannose residues with mannosidase I;  $\text{Man}_5\text{GlcNAc}_2$  is further modified  
5 by adding GlcNAc and removing 2 more mannose residues, followed by sequentially adding GlcNAc, galactose (Gal), and N-acetylneuraminic acid (also called sialic acid (NeuNAc)) to form various hybrid or complex sugar chains (R. Kornfeld and S. Kornfeld, *Ann. Rev. Biochem.* 54: 631-664, 1985; Chiba et al *J. Biol. Chem.* 273: 26298-26304, 1998).

10 In yeast, the modification of the sugar chain in the Golgi involves a series of additions of mannose residues by different mannosyltransferases ("outer chain" glycosylation). The structure of the outer chain glycosylation is specific to the organisms, typically with more than 50 mannose residues in *S. cerevisiae*, and most commonly with structures smaller than  $\text{Man}_{14}\text{GlcNAc}_2$  in *Pichia pastoris*. This yeast-specific outer chain glycosylation of the high mannose type is also denoted  
15 hyperglycosylation.

Hyperglycosylation is often undesired since it leads to heterogeneity of a recombinant protein product in both carbohydrate composition and molecular weight, which may complicate the protein purification. The specific activity (units/weight) of  
20 hyperglycosylated enzymes may be lowered by the increased portion of carbohydrate. In addition, the outer chain glycosylation is strongly immunogenic which is undesirable in a therapeutic application. Moreover, the large outer chain sugar can mask the immunogenic determinants of a therapeutic protein. For example, the influenza neuraminidase (NA) expressed in *P. pastoris* is glycosylated with N-glycans  
25 containing up to 30-40 mannose residues. The hyperglycosylated NA has a reduced immunogenicity in mice, as the variable and immunodominant surface loops on top of the NA molecule are masked by the N-glycans (Martinet et al. *Eur J. Biochem.* 247: 332-338, 1997).

Therefore, it is desirable to genetically engineer methylotrophic yeast strains in which glycosylation of proteins can be manipulated and from which recombinant proteins can be produced that would not be compromised in structure or function by large N-glycan side chains.

5

### Summary of the Invention

The present invention is directed to methods and vectors useful for genetically modifying the glycosylation process in methylotrophic yeast strains to produce glycoproteins with reduced glycosylation. Methylotrophic yeast strains  
10 generated using the present methods and vectors, as well as glycoproteins produced from such genetically modified strains are also provided.

In one embodiment, the present invention provides vectors useful for making genetically engineered methylotrophic yeast strains which are capable of producing glycoproteins with reduced glycosylation.

15 In one aspect, the present invention provides "knock-in" vectors which are capable of expressing in a methylotrophic yeast strain one or more proteins whose enzymatic activities lead to a reduction of glycosylation in glycoproteins produced by the methylotrophic yeast strain.

In a preferred embodiment, the knock-in vectors of the present invention  
20 include a nucleotide sequence coding for an  $\alpha$ -1,2-mannosidase or a functional part thereof and are capable of expressing the  $\alpha$ -1,2-mannosidase or the functional part in a methylotrophic yeast strain. A preferred nucleotide sequence is a nucleotide sequence encoding the  $\alpha$ -1,2-mannosidase of a fungal species, and more preferably, *Trichoderma reesei*. Preferably, the  $\alpha$ -1,2-mannosidase expression vector is  
25 engineered such that the  $\alpha$ -1,2-mannosidase or a functional part thereof expressed from the vector includes an ER-retention signal. A preferred ER-retention signal is HDEL. The  $\alpha$ -1,2-mannosidase coding sequence can be operable linked to a constitutive or inducible promoter, and a 3' termination sequence. The vectors can be

integrative vectors or replicative vectors. Particularly preferred  $\alpha$ -1,2-mannosidase expression vectors include pGAPZMFMAnHDEL, pGAPZMFMAnMycHDEL, pPICZBMFMAnMycHDEL, pGAPZmManHDEL, pGAPZmMycManHDEL, pPIC9mMycManHDEL and pGAPZmMycManHDEL.

- 5 In another preferred embodiment, the knock-in vectors of the present invention include a sequence coding for a glucosidase II or a functional part thereof and are capable of expressing the glucosidase II or the functional part in a methylotrophic yeast strain. A preferred nucleotide sequence is a nucleotide sequence encoding the glucosidase II of a fungal species, and more preferably, *Saccharomyces*
- 10 *cerevisiae*. Preferably, the glucosidase II expression vector is engineered such that the glucosidase II or a functional part thereof expressed from the vector includes an ER-retention signal. A preferred ER-retention signal is HDEL. The glucosidase II coding sequence can be operable linked to a constitutive or inducible promoter, and a 3' termination sequence. The vectors can be integrative vectors or replicative vectors.
- 15 Particularly preferred glucosidase II expression vectors include pGAPZAGLSII, pPICZAGLSII, pAOX2ZAGLSII, pYPTIZAGLSII, pGAPADEglsII, pPICADEglsII, pAOX2ADEglsII, pYPTIADEglsII, pGAPZAglslIHDEL and pGAPADEglsIIHDEL.

Expression vectors which include both of an  $\alpha$ -1,2-mannosidase expression unit and a glucosidase II expression unit are also provided by the present invention.

- 20 In another aspect, the present invention provides "knock-out" vectors which, when introduced into a methylotrophic yeast strain, inactivate or disrupt a gene thereby facilitating the reduction in the glycosylation of glycoproteins produced in the methylotrophic yeast strain.

- In one embodiment, the present invention provides a "knock-out" vector
- 25 which, when introduced into a methylotrophic yeast strain, inactivates or disrupts the Och1 gene. A preferred Och1 knock-out vector is pBLURA5'PpOCH1.

Still another embodiment of the present invention provides vectors which include both a knock-in unit and a knock-out unit.

Furthermore, any of the knock-in or knock-out vectors of the present invention can also include a nucleotide sequence capable of expressing a heterologous protein of interest in a methylotrophic yeast.

Another embodiment of the present invention provides methods of  
5 modifying the glycosylation in a methylotrophic yeast by transforming the yeast with one or more vectors of the present invention.

Strains of a methylotrophic yeast which can be modified using the present methods include, but are not limited to, yeast strains capable of growth on methanol such as yeasts of the genera *Candida*, *Hansenula*, *Torulopsis*, and *Pichia*. Preferred  
10 methylotrophic yeasts are of the genus *Pichia*. Especially preferred are *Pichia pastoris* strains GS115 (NRRL Y-15851), GS190 (NRRL Y-18014), PPF1 (NRRL Y-18017), PPY120H, yGC4, and strains derived therefrom. Methylotrophic yeast strains which can be modified using the present methods also include those methylotrophic yeast strains which have been engineered to express one or more heterologous proteins  
15 of interest. The glycosylation on the heterologous proteins expressed from these previously genetically engineered strains can be reduced by transforming such strains with one or more of the vectors of the present invention

Methylotrophic yeast strains which are modified by practicing the present methods are provided in another embodiment of the present invention.

20 A further aspect of the present invention is directed to methods of producing glycoproteins with a reduced glycosylation.

In accordance with such methods, a nucleotide sequence capable of expressing a glycoprotein can be introduced into a methylotrophic yeast strain which has previously been transformed with one or more of the vectors of the present  
25 invention. Alternatively, a methylotrophic yeast strain which has been genetically engineered to express a glycoprotein can be transformed with one or more of the vectors of the present invention. Moreover, if a methylotrophic yeast strain is not transformed with a nucleotide sequence encoding a glycoprotein of interest or any of the vectors of the present invention, such yeast strain can be transformed, either

consecutively or simultaneously, with both a nucleotide sequence capable of expressing the glycoprotein and one or more vectors of the present invention.

Additionally, a methylotrophic yeast strain can be transformed with one or more of the present knock-in and/or knock-out vectors which also include a nucleotide sequence  
5 capable of expressing a glycoprotein in the methylotrophic yeast strain.

Glycoproteins products produced by using the methods of the present invention, i.e., glycoproteins with reduced N-glycosylation, are also part of the present invention.

Kits which include one or more of the vectors of the present invention, or  
10 one or more strains modified to produce glycoproteins with reduced glycosylation, are also provided.

#### Brief Description of the Drawings

**Figure 1** depicts vectors carrying an HDEL-tagged *Trichoderma reesei*  $\alpha$ -  
15 1,2-mannosidase expression cassette and describes the way in which these vectors were constructed according to methods known in the art. Abbreviations used throughout construction schemes: 5' AOX1 or AOX1 P: *Pichia pastoris* AOX1 promoter sequence; Amp R: ampicillin resistance gene; ColE1: ColE1 origin of replication; 3'AOX1: 3' sequences of the *Pichia pastoris* AOX1 gene; HIS4: HIS4  
20 gene of *Pichia pastoris*. AOX TT: transcription terminator sequence of the *Pichia pastoris* AOX1 gene; ORF: open reading frame; S: secretion signal; P TEF1: the promoter sequence of the *Saccharomyces cerevisiae* transcription elongation factor 1 gene; P EM7: synthetic constitutive prokaryotic promoter EM7; Zeocin: Zeocin resistance gene; CYC1 TT: 3' end of the *S. cerevisiae* CYC1 gene; GAP: promoter  
25 sequence of the *Pichia pastoris* glyceraldehyde-3-phosphate dehydrogenase gene; PpURA3: *Pichia pastoris* URA3 gene. As can be seen in this figure, the *Trichoderma reesei*  $\alpha$ -1,2-mannosidase was operably linked to the coding sequence for the *S. cerevisiae*  $\alpha$ -mating factor secretion signal sequence and further operably linked at the 3' terminus of the coding sequence to the coding sequence for an HDEL peptide. The

whole fusion construct was operably linked to either the *P. pastoris* AOX1 promoter (in pPIC9MFManHDEL) or to the *P. pastoris* GAP promotor (in pGAPZMFManHDEL).

**Figure 2** depicts vectors carrying an HDEL-tagged *Mus musculus*  $\alpha$ -1,2-mannosidase IB expression cassette and describes the way in which these vectors were constructed according to methods known in the art. As can be seen in this figure, the catalytic domain of the *Mus musculus*  $\alpha$ -1,2-mannosidase IB was operably linked to the coding sequence for the *S. cerevisiae*  $\alpha$ -mating factor secretion signal sequence and further operably linked at the 3' terminus of the coding sequence to the coding sequence for an HDEL peptide. The whole fusion construct was operably linked to either the *P. pastoris* AOX1 promoter (in pPIC9mManHDEL) or to the *P. pastoris* GAP promotor (in pGAPZmManHDEL). Furthermore, variants of the expression cassette were made in which the coding sequence for a cMyc epitope tag was inserted between the coding sequence for the *S. cerevisiae*  $\alpha$ -Mating Factor secretion signal sequence and the coding sequence for the catalytic domain of the *Mus musculus*  $\alpha$ -1,2-mannosidase IB. This expression cassette was also operably linked to either the *P. pastoris* AOX1 promoter (in pPIC9mMycManHDEL) or to the *P. pastoris* GAP promotor (in pGAPZmMycManHDEL).

**Figure 3** depicts vectors carrying a MycHDEL tagged *Trichoderma reesei*  $\alpha$ -1,2-mannosidase and the way in which these vectors were obtained. The resulting fusion construction was again operably linked to either the *P. pastoris* AOX1 promoter (in pPICZBMFManMycHDEL) or to the *P. pastoris* GAP promotor (in pGAPZMFManMycHDEL).

**Figure 4** demonstrates the intracellular localization of the MycHDEL-tagged *Trichoderma reesei*  $\alpha$ -1,2-mannosidase and indicates ER-targeting by immunofluorescence analysis. **Panel A** Western blotting. Yeast strains were grown in 10 ml YPG cultures to an  $OD_{600}=10$ , diluted fivefold and grown in YPM for 48 h. 1/50th of the culture medium and 1/65th of the cells were analysed by SDS-PAGE and



Western blotting with the mouse monoclonal 9E10 anti-Myc antibody. The position of molecular weight marker proteins are indicated with arrows. Lanes 1-5: cellular lysates. 1,2: pGAPZMFManMycHDEL transformants. 3: non-transformed PPY12OH (negative control). 4,5: pPICZBMFManMycHDEL transformants. Lanes 6-10: culture media. 6: non transformed PPY12OH (negative control). 7,8: pGAPZMFManMycHDEL transformants. 9,10: pPICZBMFManMycHDEL transformants. **Panel B** Immunofluorescence microscopy. 1: phase contrast image of a *P. pastoris* cell (strain PPY12OH transformed with pGAPZMFManHDEL) at 1000x magnification. The nucleus is visible as an ellipse in the lower right quadrant of the cell. 2: same cell as in 1, but in fluorescence microscopy mode to show localization of the *T. reesei* mannosidase-Myc-HDEL protein. The protein is mainly localized in a circular distribution around the nucleus (nuclear envelope), which is typical for an endoplasmic reticulum steady-state distribution. 3: phase contrast image of a *P. pastoris* cell (strain PPY12OH transformed with pGAPZMFManHDEL) at 1000x magnification. 4: same cell in fluorescence microscopy to show localization of the Golgi marker protein OCH1-HA in *P. pastoris* strain PPY12OH. The dot-like distribution throughout the cytoplasm, with 3-4 dots per cell is typical for cis-Golgi distribution in *P. pastoris*.

**Figure 5** depicts the co-sedimentation of mannosidase-MycHDEL with Protein Disulfide Isomerase in sucrose density gradient centrifugation. The top panel shows the distribution over the different fractions of the sucrose gradient of the OCH1-HA Golgi marker protein. The middle panel shows this distribution for the Protein Disulfide Isomerase endoplasmic reticulum marker protein. Finally, the bottom panel shows the distribution of the MycHDEL-tagged *Trichoderma reesei*  $\alpha$ -1,2-mannosidase over the same fractions. It is concluded that the mannosidase-MycHDEL almost exactly matches the distribution of the ER marker PDI and thus mainly resides in the ER of the *Pichia pastoris* yeast cells.

**Figure 6** depicts the N-glycan analysis of *Trypanosoma cruzi* trans-sialidase coexpressed with *Trichoderma reesei* mannosidase-HDEL. Panel A: malto-

oligosaccharide size reference ladder. Sizes of the glycans are expressed in Glucose Units (GU) by comparison of their electrophoretic mobility to the mobility of these malto-oligosaccharides. Panel B: N-glycans derived from recombinant *Trypanosoma cruzi* trans-sialidase expressed in *Pichia pastoris*. The peak at GU=9,2 corresponds to

5 Man<sub>8</sub>GlcNAc<sub>2</sub>. Panel C: same analytes as panel 2, but after overnight treatment with 3U/ml purified recombinant *T. reesei* α-1,2-mannosidase. Panel D: N-glycans derived from recombinant trans-sialidase co-expressed in *Pichia pastoris* with *T. reesei* mannosidase-HDEL (under control of the GAP promotor). The peak at GU=7,6 corresponds to the Man<sub>5</sub>GlcNAc<sub>2</sub> peak in the profile of RNase B (Panel F).

10 Panel E: same analytes as panel D, but after overnight treatment with 3 mU/ml purified recombinant *T. reesei* α-1,2-mannosidase. Panel F: N-glycans derived from bovine RNase B. These glycans consist of Man<sub>5</sub>GlcNAc<sub>2</sub> to Man<sub>8</sub>GlcNAc<sub>2</sub>. Different isomers are resolved, accounting for the number of peaks for Man<sub>7</sub>GlcNAc<sub>2</sub>.

**Figure 7** depicts the processing of influenza haemagglutinin N-glycans by

15 HDEL-tagged *Trichoderma reesei* α-1,2-mannosidase and the HDEL-tagged catalytic domain of murine α-1,2-mannosidase IB. The Man<sub>5</sub>GlcNAc<sub>2</sub> reference oligosaccharide runs at scan 1850 in this analysis (not shown). Panel 1: malto-oligosaccharide size reference ladder. Panel 2: N-glycans derived from recombinant influenza haemagglutinin expressed in *Pichia pastoris*. The peak at scan 2250

20 corresponds to Man<sub>9</sub>GlcNAc<sub>2</sub>. Panel 3: N-glycans derived from recombinant haemagglutinin co-expressed in *Pichia pastoris* with *T. reesei* mannosidase-HDEL (under control of the GAP promotor). The peak at scan 1950 corresponds to Man<sub>6</sub>GlcNAc<sub>2</sub>. Panel 4: Same analytes as for panel 3, but after overnight treatment with 30 mU purified recombinant *T. reesei* α-1,2-mannosidase. Panel 5: N-glycans

25 derived from recombinant haemagglutinin co-expressed in *Pichia pastoris* with mouse mannosidase IB-HDEL (under control of the GAP promotor). Panel 6: same analytes as for panel 5, but after overnight treatment with 30 mU purified recombinant *T. reesei* α-1,2-mannosidase.

**Figure 8** graphically depicts vector pBLURA5'PpOCH1 and the way in which it was constructed.

**Figure 9** depicts the scheme for disrupting the *Pichia pastoris* OCH1 gene by single homologous recombination using pBLURA5'PpOCH1.

5       **Figure 10** depicts the cell wall glycoprotein N-glycan analysis of the Och1-inactivated clone and three clones derived from this Och1-inactivated clone by transformation with pGAPZMFManHDEL. Panel 1 shows the analysis of a mixture of malto-oligosaccharides, the degree of polymerisation of which is given by the numbers on the very top of the figure. This analysis serves as a size reference for the  
10 other panels. On the vertical axis of all panels, peak intensity in relative fluorescence units is given. Panel 2-6: analysis of the cell wall glycoprotein N-glycans of the following strains: Panel 2, non-engineered *P. pastoris* strain yGC4; Panel 3, yGC4 transformed with pBLURA5'PpOch1; 4-6, three clones of the strain of Panel 3, supplementarily transformed with pGAPZMFManHDEL. Panel 7: the N-glycans  
15 derived from bovine RNaseB, consisting of a mixture of Man<sub>5-9</sub>GlcNAc<sub>2</sub>. As can be seen from comparison between panel 2 and 3 and reference to panel 7, transformation with pBLURA5'PpOch1 leads to a strongly increased abundance of the Man<sub>8</sub>GlcNAc<sub>2</sub> substrate N-glycan (named peak 1 in Panel 2) of OCH1p. Peak 2 represents the Man<sub>9</sub>GlcNAc<sub>2</sub> product of OCH1p. Furthermore, upon supplementary transformation  
20 of pGAPZMFManHDEL, the major glycan on the cell wall glycoproteins of three independent clones is the Man<sub>5</sub>GlcNAc<sub>2</sub> end product (peak 3 in panel 4) of *T. reesei*  $\alpha$ -1,2-mannosidase digestion of the Man<sub>8</sub>GlcNAc<sub>2</sub> substrate.

**Figure 11** depicts the analysis of exactly the same glycan mixtures as in Figure 10, but after an *in vitro* digest with 3mU/ml purified *Trichoderma reesei*  $\alpha$ -1,2-mannosidase, overnight in 20 mM sodium acetate pH=5.0. Axis assignment is the  
25 same as in Figure 10. More Man<sub>5</sub>GlcNAc<sub>2</sub> is formed in the pBLURA5'PpOch1 transformed strain (Panel 3) than in the parent strain (Panel 2). Peaks in all panels before scan 3900 come from contaminants and should be ignored in the analysis.

**Figure 12** depicts the expression vector pGAPZAGLSII (SEQ ID NO: 18).  
 P TEF1: promotor of *S. cerevisiae* transcription elongation factor gene. P Em7:  
 synthetic prokaryotic promotor. Zeocin: zeocine resistance marker gene. CYC1 TT:  
 transcription terminator of *S. cerevisiae* cytochrome C1 gene. Co1 E1: bacterial origin  
 5 of replication. GAP: promotor of the *P. pasttoris* GAP gene. GLS2: *S. cerevisiae*  
 glucosidase II gene. AOX1 TT: transcription terminator of the *P. pastoris* AOX1 gene

**Figure 13** depicts the expression vector pAOX2ZAGLSII (SEQ ID NO:  
 16). P TEF1: promotor of *S. cerevisiae* transcription elongation factor gene. P Em7:  
 10 synthetic prokaryotic promotor. Zeocin: zeocine resistance marker gene. CYC1 TT:  
 transcription terminator of *S. cerevisiae* cytochrome C1 gene. Co1 E1: bacterial origin  
 of replication. AOX2 P: promotor of the *P. pastoris* AOX2 gene. GLS2: *S. cerevisiae*  
 glucosidase II gene. AOX1 TT: transcription terminator of the *P. pastoris* AOX1 gene

**Figure 14** depicts the expression vector pPICZAGLSII (SEQ ID NO: 20).  
 15 P TEF1: promotor of *S. cerevisiae* transcription elongation factor gene. P Em7:  
 synthetic prokaryotic promotor. Zeocin: zeocine resistance marker gene. CYC1 TT:  
 transcription terminator of *S. cerevisiae* cytochrome C1 gene. Co1 E1: origin of  
 replication. AOX1 P: promotor of the *P. pastoris* AOX1 gene. GLS2: *S. cerevisiae*  
 glucosidase II gene. AOX1 TT: transcription terminator of the *P. pastoris* AOX1 gene

**Figure 15** depicts the expression vector pYPT1ZAGLSII ((SEQ ID NO:  
 22). P TEF1: promotor of *S. cerevisiae* transcription elongation factor gene. P Em7:  
 synthetic prokaryotic promotor. Zeocin: zeocine resistance marker gene. CYC1 TT:  
 transcription terminator of *S. cerevisiae* cytochrome C1 gene. Co1 E1: origin of  
 replication. P YPT1: promotor of the *P. pastoris* YPT1 gene. GLS2: *S. cerevisiae*  
 25 glucosidase II gene. AOX1 TT: transcription terminator of the *P. pastoris* AOX1  
 gene.

**Figure 16** depicts the expression vector pGAPADE1glsII (SEQ ID NO:  
 19). Amp R: Ampillicin resistance marker gene. ADE1: *P. pastoris* ADE1 selection

marker gene. GAP: promotor of the P. Pastoris GAP gene. GLS2: S. cerevisiae glucosidase II gene. AOX1 TT: transcription terminator of the P. pastoris AOX1 gene

**Figure 17** depicts the expression vector pAOX2ADE1glsII (SEQ ID NO: 17). Amp R: Ampillicin resistance marker gene. ADE1: P. pastoris ADE1 selection marker gene. AOX2 P: promotor of the P. pastoris AOX2 gene. GLS2: S. cerevisiae glucosidase II gene. AOX1 TT: transcription terminator of the P. pastoris AOX1 gene.

**Figure 18** depicts the expression vector pPICADE1glsII (SEQ ID NO: 21). Amp R: Ampillicin resistance marker gene. ADE1: P. pastoris ADE1 selection marker gene. AOX1 P: promotor of the P. pastoris AOX1 gene. GLS2: S. cerevisiae glucosidase II gene. AOX1 TT: transcription terminator of the P. pastoris AOX1 gene.

**Figure 19** depicts the expression vector pYPT1ADE1glsII (SEQ ID NO: 23). Amp R: Ampillicin resistance marker gene. ADE1: P. pastoris ADE1 selection marker gene. P YPT1: promotor of the P. pastoris YPT1 gene. GLS2: S. cerevisiae glucosidase II gene. AOX1 TT: transcription terminator of the P. pastoris AOX1 gene.

**Figure 20** depicts the expression vector pGAPZAglslIHDEL (SEQ ID NO: 24). Amp R: Ampillicin resistance marker gene. ADE1: P. pastoris ADE1 selection marker gene. GAP: promotor of the P. pastoris GAP gene. GLS2: S. cerevisiae glucosidase II gene. AOX1 TT: transcription terminator of the P. pastoris AOX1 gene.

**Figure 21** depicts the expression vector pGAPADE1glsIIHDEL (SEQ ID NO: 25). P TEF1: promotor of S. cerevisiae transcription elongation factor gene. P Em7: synthetic prokaryotic promotor. Zeocin: zeocine resistance marker gene. CYC1 TT: transcription terminator of S. cerevisiae cytochrome C1 gene. Col E1: bacterial origin of replication. GAP: promotor of the P. pastoris GAP gene. GLS2: S. cerevisiae glucosidase II gene. AOX1 TT: transcription terminator of the P. pastoris AOX1 gene.

**Figure 22** depicts the test of the GLSII activity assay using a commercially available yeast alpha-glucosidase (Sigma: Cat. No. G-5003). The assay mixture contains phosphate-citrate buffer pH 6.8, mannose, 2-deoxy-D-glucose, the substrate 4-methylumbelliferyl-alpha-D-glucopyranoside and alpha-glucosidase from Sigma. 1: assay mixture illuminated with UV-light after overnight incubation at 37 ° C; 2: same as 1, but this time, the assay mixture lacks the alpha-glucosidase; 3: same as 1, but this time, the assay mixture lacks the substrate.

**Figure 23** depicts the results of the activity of recombinantly expressed GLSII from *Pichia pastoris*. All assay mixtures were incubated overnight at 37 °C and afterwards illuminated with UV-light. 1: assay with yeast alpha-glucosidase (Sigma: Cat. No. G-5003); 2: assay with the purified medium of strain 18 (PPY12-OH transformed with pGAPZAGLSII); 3: assay with purified medium of the WT PPY12-OH strain; 4: assay with the purified medium of strain H3 (PPY12-OH transformed with pGAPZAglslIHDEL).

#### Detailed Description of the Invention

It has been established that the majority of N-glycans on glycoproteins leaving the endoplasmic reticulum (ER) of *Pichia* have the core Man<sub>8</sub>GlcNAc<sub>2</sub> oligosaccharide structure. After the proteins are transported from the ER to the Golgi apparatus, additional mannose residues are added to this core sugar moiety by different mannosyltransferases, resulting in glycoproteins with large mannose side chains. Such hyperglycosylation of recombinant glycoproteins is undesirable in many instances. Accordingly, the present invention provides methods and vectors for genetically modifying methylotrophic yeast strains to produce glycoproteins with reduced glycosylation. Methylotrophic yeast strains generated using the present methods and vectors, as well as glycoproteins produced from such genetically modified strains are also provided.

In one embodiment, the present invention provides vectors useful for genetically modifying methylotrophic yeast strains to produce glycoproteins with reduced glycosylation.

In one aspect, the present invention provides “knock-in” vectors which are  
5 capable of expressing in a methylotrophic yeast strain one or more proteins whose enzymatic activities lead to a reduction of glycosylation in glycoproteins produced by the methylotrophic yeast strain. According to the present invention, such proteins include, e.g., an  $\alpha$ -1,2-mannosidase, a glucosidase II, or functional parts thereof.

In a preferred embodiment, the vectors of the present invention include a  
10 sequence coding for an  $\alpha$ -1,2-mannosidase or a functional part thereof and are capable of expressing the  $\alpha$ -1,2-mannosidase or the functional part in a methylotrophic yeast strain.

An  $\alpha$ -1,2-mannosidase cleaves the  $\alpha$ -1,2-linked mannose residues at the non-reducing ends of  $\text{Man}_8\text{GlcNAc}_2$ , and converts this core oligosaccharide on  
15 glycoproteins to  $\text{Man}_5\text{GlcNAc}_2$ . *In vitro*,  $\text{Man}_5\text{GlcNAc}_2$  is a very poor substrate for any *Pichia* Golgi mannosyltransferase, i.e., mannose residues can not be added to this sugar structure. On the other hand,  $\text{Man}_5\text{GlcNAc}_2$  is the acceptor substrate for the mammalian N-acetylglucosaminyl-transferase I and is an intermediate for the hybrid- and complex-type sugar chains characteristic of mammalian glycoproteins. Thus, by  
20 way of introducing an  $\alpha$ -1,2-mannosidase into methylotrophic yeasts such as *Pichia*, glycoproteins with reduced mannose content can be produced.

According to the present invention, the nucleotide sequence encoding an  $\alpha$ -1,2-mannosidase for use in the expression vector of the present invention can derive from any species. A number of  $\alpha$ -1,2-mannosidase genes have been cloned and are  
25 available to those skilled in the art, including mammalian genes encoding, e.g., a murine  $\alpha$ -1,2-mannosidase (Herscovics et al. *J. Biol. Chem.* 269: 9864-9871, 1994), a rabbit  $\alpha$ -1,2-mannosidase (Lal et al. *J. Biol. Chem.* 269: 9872-9881, 1994) or a human  $\alpha$ -1,2-mannosidase (Tremblay et al. *Glycobiology* 8: 585-595, 1998), as well as fungal

genes encoding, e.g., an *Aspergillus*  $\alpha$ -1,2-mannosidase (msdS gene), a *Trichoderma reesei*  $\alpha$ -1,2-mannosidase (Maras et al. *J. Biotechnol.* 77: 255-263, 2000), or a *Saccharomyces cerevisiae*  $\alpha$ -1,2-mannosidase. Protein sequence analysis has revealed a high degree of conservation among the eukaryotic  $\alpha$ -1,2-mannosidases identified so far.

Preferably, the nucleotide sequence for use in the present vectors encodes a fungal  $\alpha$ -1,2-mannosidase, more preferably, a *Trichoderma reesei*  $\alpha$ -1,2-mannosidase, and more particularly, the *Trichoderma reesei*  $\alpha$ -1,2-mannosidase described by Maras et al. *J. Biotechnol.* 77: 255-63 (2000).

According to the present invention, the nucleotide sequence can also code for only a functional part of an  $\alpha$ -1,2-mannosidase.

By "functional part" is meant a polypeptide fragment of an  $\alpha$ -1,2-mannosidase which substantially retains the enzymatic activity of the full-length protein. By "substantially" is meant at least about 40%, or preferably, at least 50% or more of the enzymatic activity of the full-length  $\alpha$ -1,2-mannosidase is retained. For example, as illustrated by the present invention, the catalytic domain of the murine  $\alpha$ -1,2-mannosidase IB constitutes a "functional part" of the murine  $\alpha$ -1,2-mannosidase IB. Those skilled in the art can readily identify and make functional parts of an  $\alpha$ -1,2-mannosidase using a combination of techniques known in the art. Predictions of the portions of an  $\alpha$ -1,2-mannosidase essential to or sufficient to confer the enzymatic activity can be made based on analysis of the protein sequence. The activity of a portion of an  $\alpha$ -1,2-mannosidase of interest, expressed and purified from an appropriate expression system, can be verified using *in vitro* or *in vivo* assays described hereinbelow.

In accordance with the present invention, an  $\alpha$ -1,2-mannosidase or a functional part thereof expressed in a methylotrophic yeast strain preferably is targeted to a site in the secretory pathway where  $\text{Man}_8\text{GlcNAc}_2$  (the substrate of  $\alpha$ -1,2-



mannosidase) is already formed on a glycoprotein, but has not reached a Golgi glycosyltransferase which elongates the sugar chain with additional mannose residues.

Accordingly, in a preferred embodiment of the present invention, the  $\alpha$ -1,2-mannosidase expression vector is engineered as such that the  $\alpha$ -1,2-mannosidase  
5 or a functional part thereof expressed from the vector includes an ER-retention signal.

"An ER retention signal" refers to a peptide sequence which directs a protein having such peptide sequence to be transported to and retained in the ER. Such ER retention sequences are often found in proteins that reside and function in the ER.

10 Multiple choices of ER retention signals are available to those skilled in the art, e.g., the first 21 amino acid residues of the *S. cerevisiae* ER protein MNS1 (Martinet et al. *Biotechnology Letters* 20: 1171-1177, 1998). A preferred ER retention signal for use in the present invention is peptide HDEL (SEQ ID NO: 1). The HDEL peptide sequence, found in the C-terminus of a number of yeast proteins, acts as a  
15 retention/retrieval signal for the ER (Pelham *EMBO J.* 7: 913-918, 1988). Proteins with an HDEL sequence are bound by a membrane-bound receptor (Erd2p) and then enter a retrograde transport pathway for return to the ER from the Golgi apparatus.

According to the present invention, an ER retention signal can be placed anywhere in the protein sequence of an  $\alpha$ -1,2-mannosidase, but preferably at the C-  
20 terminus of the  $\alpha$ -1,2-mannosidase.

The  $\alpha$ -1,2-mannosidase for use in the present invention can be further modified, e.g., by insertion of an epitope tag to which antibodies are available, such as Myc, HA, FLAG and His6 tags well-known in the art. An epitope-tagged  $\alpha$ -1,2-mannosidase can be conveniently purified, or monitored for both expression and  
25 intracellular localization.

An ER retention signal and an epitope tag can be readily introduced into a protein of interest by inserting nucleotide sequences coding for such signal or tag into

the nucleotide sequence encoding the protein of interest, using any of the molecular biology techniques known in the art.

In another preferred embodiment, the vectors of the present invention include a sequence coding for a glucosidase II or a functional part thereof and are  
5 capable of expressing the glucosidase II or the functional part in the methylotrophic yeast strain.

It has been established that the initial N-linked oligosaccharide (Glc<sub>3</sub>Man<sub>9</sub>GlcNAc<sub>2</sub>), transferred in the ER onto a protein, is cleaved in the ER by specific glucosidases to remove the glucose residues, and by a mannosidase to remove  
10 one specific  $\alpha$ -1,2-linked mannose. It has been observed by the present inventors that some recombinant proteins expressed in *Pichia* have residual glucose residues on the sugar moiety when such proteins leave the ER for the Golgi apparatus. The residual glucose molecules present on the sugar structure prevent the complete digestion of the sugar moiety by an  $\alpha$ -1,2-mannosidase, and the introduction of an exogenous  
15 glucosidase can facilitate the removal of these glucose residues.

According to the present invention, the nucleotide sequence encoding a glucosidase II can derive from any species. Glucosidase II genes have been cloned from a number of mammalian species including rat, mouse, pig and human. The glucosidase II protein from these mammalian species consists of an alpha and a beta  
20 subunit. The alpha subunit is about 110 kDa and contains the catalytic activity of the enzyme, while the beta subunit has a C-terminal HDEL ER-retention sequence and is believed to be important for the ER localization of the enzyme. The glucosidase II gene from *S. cerevisiae* has also been cloned (ORF YBR229c, located on chromosome II). This gene encodes a protein of about 110 kDa, which shows a high degree of  
25 homology to the mammalian alpha subunits.

A preferred glucosidase II gene for use in the present vectors is from a fungal species such as *Pichia pastoris* and *S. cerevisiae*. An example of a fungal glucosidase II gene is the *S. cerevisiae* glucosidase II alpha subunit gene.

According to the present invention, the nucleotide sequence can also encode only a functional part of a glucosidase II. By "functional part" is meant a polypeptide fragment of a glucosidase II which substantially retains the enzymatic activity of the full-length protein. By "substantially" is meant at least about 40%, or  
5 preferably, at least 50% or more of the enzymatic activity of the full-length glucosidase II is retained. Functional parts of a glucosidase II can be identified and made by those skilled in the art using a variety of techniques known in the art.

In a preferred embodiment of the present invention, the glucosidase II protein is engineered to include an ER retention signal such that the protein expressed  
10 in a methylotrophic yeast strain is targeted to the ER and retains therein for function. ER retention signals are as described hereinabove, e.g., the HDEL peptide sequence.

The glucosidase II for use in the present invention can be further modified, e.g., by insertion of an epitope tag to which antibodies are available, such as Myc, HA, FLAG, and His6 tag, which are well-known in the art.

15 According to the present invention, the "knock-in" vectors can include either or both of an  $\alpha$ -1,2-mannosidase coding sequence and a glucosidase II coding sequence.

Further according to the present invention, the nucleotide sequence coding for the enzyme to be expressed (e.g., an  $\alpha$ -1,2-mannosidase or a functional part  
20 thereof, or a glucosidase II or a functional part thereof) can be placed in an operable linkage to a promoter and a 3' termination sequence.

Promoters appropriate for expression of a protein in a methylotrophic yeast can include both constitutive promoters and inducible promoters. Constitutive promoters include e.g., the *Pichia pastoris* glyceraldehyde-3-phosphate  
25 dehydrogenase promoter ("the GAP promoter"). Examples of inducible promoters include, e.g., the *Pichia pastoris* alcohol oxidase I promoter ("the AOXI promoter") (U.S. Patent No. 4,855,231), or the *Pichia pastoris* formaldehyde dehydrogenase promoter ("the FLD promoter") (Shen et al. Gene 216: 93-102, 1998).

3' termination sequences are sequences 3' to the stop codon of a structural gene which function to stabilize the mRNA transcription product of the gene to which the sequence is operably linked, such as sequences which elicit polyadenylation. 3' termination sequences can be obtained from *Pichia* or other methylotrophic yeast.

5 Examples of *Pichia pastoris* 3' termination sequences useful for the practice of the present invention include termination sequences from the *AOX1* gene, *p40* gene, *HIS4* gene and *FLD1* gene.

The vectors of the present invention preferably contain a selectable marker gene. The selectable marker may be any gene which confers a selectable phenotype  
10 upon a methylotrophic yeast strain and allows transformed cells to be identified and selected from untransformed cells. The selectable marker system may include an auxotrophic mutant methylotrophic yeast strain and a wild type gene which complements the host's defect. Examples of such systems include the *Saccharomyces cerevisiae* or *Pichia pastoris HIS4* gene which may be used to complement *his4*  
15 *Pichia* strains, or the *S. cerevisiae* or *Pichia pastoris ARG4* gene which may be used to complement *Pichia pastoris arg* mutants. Other selectable marker genes which function in *Pichia pastoris* include the *Zeo<sup>R</sup>* gene, the *G418<sup>R</sup>* gene, and the like.

The vectors of the present invention can also include an autonomous replication sequence (ARS). For example, U.S. Patent No. 4,837,148 describes  
20 autonomous replication sequences which provide a suitable means for maintaining plasmids in *Pichia pastoris*. The disclosure of U.S. Patent No. 4,837,148 is incorporated herein by reference.

The vectors can also contain selectable marker genes which function in bacteria, as well as sequences responsible for replication and extrachromosomal  
25 maintenance in bacteria. Examples of bacterial selectable marker genes include ampicillin resistance (*Amp<sup>r</sup>*), tetracycline resistance (*Tet<sup>r</sup>*), neomycin resistance, hygromycin resistance, and zeocin resistance (*Zeo<sup>R</sup>*) genes.

According to the present invention, the nucleotide sequence encoding the protein to be expressed in a methylotrophic yeast can be placed in an integrative vector or a replicative vector (such as a replicating circular plasmid).

Integrative vectors are disclosed, e.g., in U.S. Patent No. 4,882,279 which  
5 is incorporated herein by reference. Integrative vectors generally include a serially arranged sequence of at least a first insertable DNA fragment, a selectable marker gene, and a second insertable DNA fragment. The first and second insertable DNA fragments are each about 200 nucleotides in length and have nucleotide sequences which are homologous to portions of the genomic DNA of the species to be  
10 transformed. A nucleotide sequence containing a structural gene of interest for expression is inserted in this vector between the first and second insertable DNA fragments whether before or after the marker gene. Integrative vectors can be linearized prior to yeast transformation to facilitate the integration of the nucleotide sequence of interest into the host cell genome.

15 Replicative and integrative vectors carrying either or both of an  $\alpha$ -1,2-mannosidase coding sequence or a glucosidase II coding sequence can be constructed by standard techniques known to one of ordinary skill in the art and found, for example, in Sambrook et al. (1989) in *Molecular Cloning: A Laboratory Manual*, or any of a myriad of laboratory manuals on recombinant DNA technology that are  
20 widely available.

Preferred vectors of the present invention carrying an  $\alpha$ -1,2-mannosidase expression sequence include pGAPZMFManHDEL, pGAPZMFManMycHDEL, pPICZBMFManMycHDEL, pGAPZmManHDEL, pGAPZmMycManHDEL, pPIC9mMycManHDEL and pGAPZmMycManHDEL, which are further described in  
25 the Examples hereinbelow.

Preferred vectors of the present invention carrying a glucosidase II expression sequence include pGAPZAGLSII, pPICZAGLSII, pAOX2ZAGLSII, pYPTIZAGLSII, pGAPADE1glsII, pPICADE1glsII, pAOX2ADE1glsII,

pYPTIADE1glsII, pGAPZAglsIIHDEL and pGAPADE1glsIIHDEL, which are further described in the Examples hereinbelow.

In another aspect, the present invention provides "knock-out" vectors which, when introduced into a methylotrophic yeast strain, inactivate or disrupt a gene thereby facilitating the reduction in the glycosylation of glycoproteins produced in the methylotrophic yeast strain.

In one embodiment, the present invention provides a "knock-out" vector which, when introduced into a methylotrophic yeast strain, inactivates or disrupts the Och1 gene.

The *S. cerevisiae* OCH1 gene has been cloned (Nakayama et al. *EMBO J.* 11: 2511-2519, 1992). It encodes a membrane bound  $\alpha$ -1,6-mannosyltransferase, localized in the early Golgi complex, that is functional in the initiation of  $\alpha$ -1,6-polymannose outer chain addition to the N-linked core oligosaccharide (Man<sub>5</sub>GlcNAc<sub>2</sub> and Man<sub>8</sub>GlcNAc<sub>2</sub>) (Nakanishi-Shindo et al. *J. Biol. Chem.* 268: 26338-26345, 1993).

A *Pichia* sequence has been described in Japanese Patent Application No. 07145005 that encodes a protein highly homologous to the *S. cerevisiae* OCH1. For purpose of the present invention, this sequence is denoted herein as "the *Pichia* OCH1 gene". Those skilled in the art can isolate the OCH1 genes from other methylotrophic yeasts using techniques well known in the art.

According to the present invention, a disruption in the OCH1 gene of a methylotrophic yeast can result in either the production of an inactive protein product or no product. The disruption may take the form of an insertion of a heterologous DNA sequence into the coding sequence and/or the deletion of some or all of the coding sequence. Gene disruptions can be generated by homologous recombination essentially as described by Rothstein (in *Methods in Enzymology*, Wu et al., eds., vol 101:202-211, 1983).

To disrupt the Och1 gene by homologous recombination, an Och1 knock-out vector can be constructed in such a way to include a selectable marker gene. The selectable marker gene is operably linked, at both 5' and 3' end, to portions of the

Och1 gene of sufficient length to mediate homologous recombination. The selectable marker can be one of any number of genes which either complement host cell auxotrophy or provide antibiotic resistance, including *URA3*, *LEU2* and *HIS3* genes. Other suitable selectable markers include the *CAT* gene, which confers  
5 chloramphenicol resistance on yeast cells, or the *lacZ* gene, which results in blue colonies due to the expression of active  $\beta$ -galactosidase. Linearized DNA fragments of an Och1 knock-out vector are then introduced into host methylotrophic yeast cells using methods well known in the art. Integration of the linear fragments into the genome and the disruption of the Och1 gene can be determined based on the selection  
10 marker and can be verified by, for example, Southern Blot analysis.

Alternatively, an Och1 knock-out vector can be constructed in such a way to include a portion of the Och1 gene to be disrupted, which portion is devoid of any Och1 promoter sequence and encodes none or an inactive fragment of the Och1 protein. By "an inactive fragment", it is meant a fragment of the Och1 protein which  
15 has, preferably, less than about 10% and most preferably, about 0% of the activity of the full-length OCH1 protein. Such portion of the OCH1 gene is inserted in a vector in such a way that no known promoter sequence is operably linked to the OCH1 sequence, but that a stop codon and a transcription termination sequence are operably linked to the portion of the Och1 gene. This vector can be subsequently linearized in  
20 the portion of the OCH1 sequence and transformed into a methylotrophic yeast strain using any of the methods known in the art. By way of single homologous recombination, this linearized vector is then integrated in the OCH1 gene. Two Och1 sequences are produced in the chromosome as a result of the single homologous recombination. The first Och1 sequence is the portion of the Och1 gene from the  
25 vector, which is now under control of the OCH1 promoter of the host methylotrophic yeast, yet cannot produce an active OCH1 protein as such Och1 sequence codes for no or an inactive fragment of the OCH1 protein, as described hereinabove. The second Och1 sequence is a full OCH1 coding sequence, but is not operably linked to any known promoter sequence and thus, no active messenger is expected to be formed for

synthesis of an active OCH1 protein. Preferably, an inactivating mutation is introduced in the OCH1 sequence, to the 5' end of the site of linearization of the vector and to the 3' end of the translation initiation codon of OCH1. By "inactivating mutation" it is meant a mutation introducing a stop codon, a frameshift mutation or any other mutation causing a disruption of the reading frame. Such mutation can be introduced into an Och1 sequence using any of the site directed mutagenesis methods known in the art. Such inactivating mutation ensures that no functional OCH1 protein can be formed even if there exist some promoter sequences 5' to the Och1 sequence in the knock-out vector.

10 A preferred Och1 knock-out vector of the present invention is pBLURA5'PpOCH1.

If desired, either or both of a mannosidase expression sequence and a glucosidase expression sequence can be carried on the same plasmid used to disrupt the OCH1 gene to create a "knock-in-and-knock-out" vector.

15 Additionally, any of the above-described vectors can further include a nucleotide sequence capable of expressing a glycoprotein of interest in a methylotrophic yeast strain.

Another aspect of the present invention is directed to methods of modifying methylotrophic yeast strains to reduce glycosylation on proteins produced by the methylotrophic yeast strains. In accordance with the present methods, methylotrophic yeast strains are modified by transforming into these yeast strains with one or more, i.e., at least one, knock-in and/or knock-out vectors of the present invention as described herein above.

25 Methylotrophic yeast strains which can be modified using the present methods include but are not limited to yeast capable of growth on methanol such as yeasts of the genera *Candida*, *Hansenula*, *Torulopsis*, and *Pichia*. A list of species which are exemplary of this class of yeasts can be found in C. Anthony (1982), *The Biochemistry of Methylotrophs*, 269. *Pichia pastoris*, *Pichia methanolica*, *Pichia anomola*, *Hansenula polymorpha* and *Candida boidinii* are examples of



methylophilic yeasts useful in the practice of the present invention. Preferred methylophilic yeasts are of the genus *Pichia*. Especially preferred are *Pichia pastoris* strains GS115 (NRRL Y-15851); GS190 (NRRL Y-18014) disclosed in U.S. Patent No. 4,818,700; PPF1 (NRRL Y-18017) disclosed in U.S. Patent No. 4,812,405; 5 PPY120H and yGC4; as well as strains derived therefrom.

Methylophilic yeast strains which can be modified using the present methods also include those methylophilic yeast strains which have been genetically engineered to express one or more heterologous glycoproteins of interest. The glycosylation on the heterologous glycoproteins expressed from these previously 10 engineered strains can be reduced by transforming such strains with one or more of the vectors of the present invention.

The vectors of the present invention can be introduced into the cells of a methylophilic yeast strain using known methods such as the spheroplast technique, described by Cregg et al. 1985, or the whole-cell lithium chloride yeast transformation 15 system, Ito et al. *Agric. Biol. Chem.* 48:341, modified for use in *Pichia* as described in EP 312,934. Other published methods useful for transformation of the plasmids or linear vectors include U.S. Patent No. 4,929,555; Hinnen et al. *Proc. Nat. Acad. Sci. USA* 75:1929 (1978); Ito et al. *J. Bacteriol.* 153:163 (1983); U.S. Patent No. 4,879,231; Sreekrishna et al. *Gene* 59:115 (1987). Electroporation and PEG1000 20 whole cell transformation procedures may also be used. Cregg and Russel *Methods in Molecular Biology: Pichia Protocols*, Chapter 3, Humana Press, Totowa, N.J., pp. 27-39 (1998).

Transformed yeast cells can be selected by using appropriate techniques including but not limited to culturing auxotrophic cells after transformation in the 25 absence of the biochemical product required (due to the cell's auxotrophy), selection for and detection of a new phenotype, or culturing in the presence of an antibiotic which is toxic to the yeast in the absence of a resistance gene contained in the transformants. Transformants can also be selected and/or verified by integration of the

expression cassette into the genome, which can be assessed by e.g., Southern Blot or PCR analysis.

In one embodiment, a methylotrophic yeast strain is transformed with a vector which includes a nucleotide sequence coding for an  $\alpha$ -1,2-mannosidase or a functional part thereof. The nucleotide sequence is capable of expressing the  $\alpha$ -1,2-mannosidase or the functional part in the methylotrophic yeast strain, and is, preferably, integrated into the genome of the methylotrophic yeast strain.

The expression of an  $\alpha$ -1,2-mannosidase introduced in a methylotrophic yeast strain can be verified both at the mRNA level, e.g., by Northern Blot analysis, and at the protein level, e.g., by Western Blot analysis. The intracellular localization of the protein can be analyzed by using a variety of techniques, including subcellular fractionation and immunofluorescence experiments. An ER localization of an  $\alpha$ -1,2-mannosidase can be determined by co-sedimentation of this enzyme with a known ER resident protein (e.g., Protein Disulfide Isomerase) in a subcellular fractionation experiment. An ER localization can also be determined by an immunofluorescence staining pattern characteristic of ER resident proteins, typically a perinuclear staining pattern.

To confirm that an  $\alpha$ -1,2-mannosidase or a functional part thereof expressed in a methylotrophic yeast strain has the expected mannose-trimming activity, both *in vitro* and *in vivo* assays can be employed. Typically, an *in vitro* assay involves digestion of an *in vitro* synthesized substrate, e.g.,  $\text{Man}_8\text{GlcNAc}_2$ , with the enzyme expressed and purified from a methylotrophic yeast strain, and assessing the ability of such enzyme to trim  $\text{Man}_8\text{GlcNAc}_2$  to, e.g.,  $\text{Man}_5\text{GlcNAc}_2$ . In *in vivo* assays, the  $\alpha$ -1,2-mannosidase or a part thereof is co-expressed in a methylotrophic yeast with a glycoprotein known to be glycosylated with N-glycans bearing terminal  $\alpha$ -1,2-linked mannose residues in such yeast. The enzymatic activity of such an  $\alpha$ -1,2-mannosidase or a part thereof can be measured based on the reduction of the number of  $\alpha$ -1,2-linked mannose residues in the structures of the N-glycans of the

glycoprotein. In both *in vitro* and *in vivo* assays, the composition of a carbohydrate group can be determined using techniques that are well known in the art and are illustrated in the Examples hereinbelow.

5 In another embodiment, a methylotrophic yeast strain is transformed with a vector which includes a nucleotide sequence coding for a glucosidase II or a functional part thereof. The nucleotide sequence is capable of expressing the glucosidase II or the functional part in the methylotrophic yeast strain, and is, preferably, integrated into the genome of the methylotrophic yeast strain.

10 The enzymatic activity of a glucosidase II or a functional part thereof expressed in a transformed methylotrophic yeast strain can be assessed using a variety of assays. For example, methylotrophic yeast cells transformed with a sequence encoding a glucosidase II or a part thereof can be set to grow on solid medium containing a substrate of the glucosidase, e.g., 5-bromo-4-chloro-3-indolyl- $\alpha$ -D-glucopyranoside or 4-MU- $\alpha$ -D-Glc. When the enzyme is expressed by the *Pichia* and  
15 secreted extracellularly, the substrate is acted upon by the enzyme, giving rise to detectable signals around the colonies such as blue color or fluorescent glow. Alternatively, liquid culture medium containing the expressed protein molecules can be collected and incubated in test tubes with a substrate, e.g., p-nitrophenyl- $\alpha$ -D-glucopyranoside. The enzymatic activity can be determined by measuring the specific  
20 product released. Moreover, *in vivo* assays can be employed, where a glucosidase II is co-expressed in yeast with a glycoprotein known to be N-glycosylated with glucose residues, e.g., influenza neuraminidase. The enzymatic activity of the glucosidase II can be measured based on the reduction of the glucose content in the sugar chain(s) of the glycoprotein.

25 In still another embodiment of the present invention, a methylotrophic yeast strain is transformed with an Och1 knock-out vector. As a result of the transformation and integration of the vector, the genomic Och1 gene in the yeast strains is disrupted.

In a further embodiment of the present invention, a methylotrophic yeast strain is transformed with any combination of an  $\alpha$ -1,2-mannosidase expression vector, a glucosidase II expression vector, and an Och1 knock-out vector. Such modification can be achieved by serial, consecutive transformations, i.e., introducing one vector at a time, or alternatively by co-transformation, i.e., introducing the vectors simultaneously.

The modified methylotrophic yeast strains described herein above can be further modified if desired. For example, additional disruption of genes encoding any other *Pichia* mannosyltransferases can be made. Genes encoding mammalian enzymes can also be introduced to produce glycoproteins having hybrid- or complex-type N-glycans, if desired.

Methylotrophic yeast strains which are modified by using the present methods, i.e., by transforming with one or more of the vectors of the present invention, form another embodiment of the present invention.

It should be understood that certain aspects of the present invention, especially the introduction of an intracellularly expressed  $\alpha$ -1,2-mannosidase activity, are also useful to obtain a reduced glycosylation of the O-linked glycans on glycoproteins produced in a methylotrophic yeast, as it is known in the art that these O-linked glycans consist mainly of  $\alpha$ -1,2-linked mannose residues. O-linked glycans as used herein refers to carbohydrate structures linked to serine or threonine residues of glycoproteins.

A further aspect of the invention is directed to methods of producing a glycoprotein with reduced glycosylation in a methylotrophic yeast, especially a glycoprotein heterologous to the methylotrophic yeast.

"A glycoprotein" as used herein refers to a protein which, in methylotrophic yeasts, is either glycosylated on one or more asparagines residues or on one or more serine or threonine residues, or on both asparagines and serine or threonine residues.

The term "reduced glycosylation" refers to a reduced size of the carbohydrate moiety on the glycoprotein, particularly with fewer mannose residues, when the glycoprotein is expressed in a methylotrophic yeast strain which has been modified in accordance with the present invention, as compared to a wild type, unmodified strain of the methylotrophic yeast.

In accordance with the present invention, the production of a glycoprotein of interest with reduced glycosylation can be achieved in a number of ways. A nucleotide sequence capable of expressing a glycoprotein can be introduced into a methylotrophic yeast strain which has been previously modified in accordance with the present invention, i.e., a strain transformed with one or more of the vectors of the present invention and capable of producing glycoproteins with reduced glycosylation. Alternatively, a methylotrophic yeast strain which has already been genetically engineered to express a glycoprotein can be transformed with one or more of the vectors of the present invention. Otherwise, if a methylotrophic yeast strain does not express a glycoprotein of interest, nor is the strain transformed with any of the vectors of the present invention, such yeast strain can be transformed, either consecutively or simultaneously, with both a nucleotide sequence capable of expressing the glycoprotein and one or more vectors of the present invention. Additionally, a methylotrophic yeast strain can be transformed with one or more of the present knock-in and/or knock-out vectors which also include a nucleotide sequence capable of expressing a glycoprotein in the methylotrophic yeast strain.

The nucleotide sequence capable of expressing a glycoprotein in a methylotrophic yeast can be made to include from 5' to 3', a promoter, a sequence encoding the glycoprotein, and a 3' termination sequence. Promoters and 3' termination sequences which are suitable for expression of a glycoprotein can include any of those promoters and 3' termination sequences described hereinabove.

The nucleotide sequence for expression of a glycoprotein can include additional sequences, e.g., signal sequences coding for transit peptides when secretion of a protein product is desired. Such sequences are widely known, readily available

and include *Saccharomyces cerevisiae* alpha mating factor prepro ( $\alpha$ mf), *Pichia pastoris* acid phosphatase (PHO1) signal sequence and the like.

The nucleotide sequence for expression of a glycoprotein can be placed on a replicative vector or an integrative vector. The choice and construction of such  
5 vectors are as described hereinabove.

The nucleotide sequence capable of expressing a glycoprotein can be carried on the same replicative plasmid as a plasmid-borne  $\alpha$ -1,2-mannosidase or glucosidase II expression unit. Alternatively, the nucleotide sequence containing the glycoprotein coding sequence is carried on a separate plasmid or integrated into the  
10 host genome.

Glycoproteins produced can be purified by conventional methods. Purification protocols can be determined by the nature of the specific protein to be purified. Such determination is within the ordinary level of skill in the art. For example, the cell culture medium is separated from the cells and the protein secreted  
15 from the cells can be isolated from the medium by routine isolation techniques such as precipitation, immunoadsorption, fractionation or a variety of chromatographic methods.

Glycoproteins which can be produced by the methods of the present invention include, e.g., *Bacillus amyloliquefaciens*  $\alpha$ -amylase, *S. cerevisiae* invertase,  
20 *Trypanosoma cruzi* trans-sialidase, HIV envelope protein, influenza virus A haemagglutinin, influenza neuraminidase, Bovine herpes virus type-1 glycoprotein D, human angiostatin, human B7-1, B7-2 and B-7 receptor CTLA-4, human tissue factor, growth factors (e.g., platelet-derived growth factor), tissue plasminogen activator, plasminogen activator inhibitor-I, urokinase, human lysosomal proteins such as  $\alpha$ -  
25 galactosidase, plasminogen, thrombin, factor XIII and immunoglobulins. For additional useful glycoproteins which can be expressed in the genetically engineered *Pichia* strains of the present invention, see Bretthauer and Castellino, *Biotechnol.*

*Appl. Biochem.* 30: 193-200 (1999), and Kukuruzinska et al. *Ann Rev. Biochem.* 56: 915-44 (1987).

Glycoproteins produced by using the methods of the present invention, i.e., glycoproteins with reduced glycosylation, are also part of the present invention.

5           Still another aspect of the present invention provides kits which contain one or more of the knock-in vectors, knock-out vectors, or knock-in-and-knock-out vectors of the present invention described above. More particularly, a kit of the present invention contains a vector capable of expressing an  $\alpha$ -mannosidase I in a methylotrophic yeast, a vector capable of expressing a glucosidase II in a  
10 methylotrophic yeast, a vector capable of disrupting the Och1 gene in a methylotrophic yeast, a vector capable of expressing both a glucosidase II and an  $\alpha$ -mannosidase, a vector a vector capable of disrupting the Och1 gene and capable of expressing either or both of a glucosidase II and an  $\alpha$ -mannosidase, or any combinations thereof.

15           The kit can also include a nucleotide sequence which encodes and is capable of expressing a heterologous glycoprotein of interest. Such nucleotide sequence can be provided in a separate vector or in the same vector which contains sequences for knocking-in or knocking out as described hereinabove.

20           In addition, the kit can include a plasmid vector in which a nucleotide sequence encoding a heterologous protein of interest can be subsequently inserted for transformation into and expression in a methylotrophic yeast. Alternatively, the knock-in or knock-out vectors in the kits have convenient cloning sites for insertion of a nucleotide sequence encoding a heterologous protein of interest.

25           The kit can also include a methylotrophic yeast strain which can be subsequently transformed with any of the knock-in, knock-out or knock-in-and-knock-out vectors described hereinabove. The kit can also include a methylotrophic yeast strain which has been transformed with one or more of the knock-in or knock-out vectors. Furthermore, the kit can include a methylotrophic yeast strain which has been

transformed with a nucleotide sequence encoding and capable of expressing a heterologous glycoprotein of interest.

The present invention is further illustrated by the following examples.



**Example 1****Introduction of  $\alpha$ -1,2-Mannosidase to the ER-Golgi Border****1.1 Plasmids**

Plasmid	Promoter	Enzyme	Tag
pGAPZMFManHDEL	GAP	<i>T. reesei</i> $\alpha$ -1,2-mannosidase	--
pGAPZMFManMycHDEL	GAP	<i>T. reesei</i> $\alpha$ -1,2-mannosidase	Myc
pPICZBMFManMycHDEL	AOX1	<i>T. reesei</i> $\alpha$ -1,2-mannosidase	Myc
pGAPZMFmManHDEL	GAP	mouse mannosidase IB catalytic domain	--
pGAPZMFmMycManHDEL	GAP	mouse mannosidase IB catalytic domain	Myc

5

The *Trichoderma reesei*  $\alpha$ -1,2-mannosidase gene has been isolated and described by Maras et al. (*J. Biotechnol.* 77;255-263, 2000). The sequence of this gene is available at NCBI Genbank under Accession No. AF212153. A construction fragment was generated by PCR using the pPIC9MFmanase plasmid (same as pPP1MFmndsl described by Maras et al. (2000)) as the template and using the following oligonucleotide primers: 5'-GACTGGTTCCAATTGACAAGC-3' (SEQ ID NO:2) and 5'-AGTCTAGATTACAACCTCGTCGTGAGCAAGGTGGCCGCCCGCCGTCG-3' (SEQ ID NO:3). The resulting product contained the 3' end of the *Pichia pastoris* AOXI promoter, the prepro-signal sequence of the *S. cerevisiae*  $\alpha$ -mating factor, the open reading frame of the *Trichoderma reesei*  $\alpha$ -1,2-mannosidase cloned in frame with the signal sequence, the coding sequence for HDEL, a stop codon and an *Xba* I restriction site. This fragment was digested with *Eco* RI and *Xba* I, removing the 5' sequences up to the mannosidase ORF, and then cloned into the vector pGAPZ $\alpha$ A (Invitrogen, Baarn, The Netherlands) which had been digested with *Eco* RI and *Xba* I, thus restoring the fusion with the *S. cerevisiae*  $\alpha$ -mating factor signal

20

sequence. The resulting plasmid was named pGAPZMFManHDEL and is graphically depicted in **Figure 1**. The ORF sequence of the MFManHDEL fusion in pGAPZMFManHDEL is set forth in SEQ ID NO: 14.

In order to introduce the coding sequence for a c-Myc tag between the catalytic domain and the HDEL-signal, the 3' end of the ORF of *T. reesei*  $\alpha$ -1,2-mannosidase was PCR-amplified using a sense primer 5'-CCATTGAGGACGCATGCCGCGCC-3' (SEQ ID NO: 4) (containing an *Sph* I restriction site) and an antisense primer GTATCTAGATTACAACTCGTCGTGCAGATCCTCTTCTGAGATGAGTTTTTGT  
 10 TCAGCAAGGTGGCCGCCCCGTCGTGATGATGAA (SEQ ID NO: 5) (containing the coding sequences of the c-Myc tag and the HDEL signal, followed by a stop codon and an *Xba* I restriction site). The resulting PCR product was digested with *Sph* I and *Xba* I, purified by agarose gel electrophoresis and inserted into pGAPZMFManHDEL which had been cut with the same restriction enzymes, resulting in plasmid  
 15 pGAPZMFManMycHDEL. To put the ORF of pGAPZMFManMycHDEL under the control of the inducible AOXI promoter, the entire ORF was liberated from pGAPZMFManMycHDEL with *Bst* BI and *Xba* I, and cloned in pPICZB (Invitrogen, Baarn, The Netherlands), resulting in pPICZBMFManMycHDEL.

Cloning of the mouse mannosidase IB catalytic domain with concomitant  
 20 addition of the coding sequence for a C-terminal HDEL-tag was done by PCR on a mouse cDNA library (mRNA isolated from the L929 cell line induced with cycloheximide and mouse Tumor Necrosis Factor. Average insert length of the cDNA library was 2000 bp). The PCR oligonucleotide primers used were:  
 25 5'AACTCGAGATGGACTCTTCAAAACACAAACGC3' (SEQ ID NO: 6) and 5'TTGGCGCCGCTTACAACTCGTCGTGTCGGACAGCAGGATTACCTGA3' (SEQ ID NO: 7). The product contained a 5' *Xho* I site and the coding sequence for C-terminal HDEL-site, followed by a stop codon and a *Not* I site at the 3' end. The product was cloned in pGAPZ $\alpha$ A via the *Xho* I /*Not* I sites in the PCR product and the vector, resulting in an in frame fusion of the mouse mannosidase catalytic domain

with the *S. cerevisiae*  $\alpha$ -mating factor signal sequence. The sequence of the entire open reading frame generated is set forth in SEQ ID NO: 15.

## **1.2 Yeast Transformation and Genomic Integration**

5

Table 2

Parental strain	DNA transformed
GS115 (his4)	pGAPZMFManHDEL
	pPIC9MFManHDEL
	pPIC9mManHDEL
	pPIC9mMycManHDEL
	pGAPZmManHDEL
	pGAPZmMycManHDEL
GS115 (his4 complemented by pPIC9InfluenzaHA)	pGAPZMFManHDEL
	pGAPZmManHDEL
	pGAPZmMycManHDEL
PPY120H (his4 complemented by pPIC9sOCH1)	pGAPZMFManMycHDEL
	pPICZBMFManMycHDEL
yGC4 (his4 arg1 ade2 ura3 complemented by pBLURA5'PpOCH1)	pPIC9InfluenzaNeuraminidase
	pGAPZMFManHDEL
	pPIC9Glucoseoxidase

All transformations to *Pichia pastoris* were performed with electroporation according to the directions of Invitrogen. Transformants of vectors carrying the Zeocin resistance gene were selected on YPD containing 100  $\mu$ g/ml Zeocine

(Invitrogen, Baarn, the Netherlands) and 1M sorbitol. Selection of transformants of pPIC9 derivatives was done on minimal medium lacking histidine and containing 1M sorbitol. Genomic integration of the expression cassettes was verified using PCR on genomic DNA purified from the *Pichia* strains using the Yeast Miniprep method (Nucleon). In all cases concerning the *Trichoderma reesei* gene fusions, the primers used were the sense primer 5'-CCATTGAGGACGCATGCCGCGCC-3' (SEQ ID NO: 8), which annealed to the 3' half of the mannosidase ORF, and the antisense primer 3' AOXI 5'-GCAAATGGCATTCTGACATCCT-3' (SEQ ID NO: 9), which annealed to the AOXI transcription terminator that was present in all our expression constructs.

For the control of genomic integration of the mouse mannosidase transgenes, PCR was done using the sense primer 5'GAP 5'GTCCCTATTTCAATCAATTGAA3' (SEQ ID NO: 10, annealing to the GAP promoter or 5'AOXI 5'GACTGGTTCCAATTGACAAGC3' (SEQ ID NO:11), annealing to AOXI promoter), and the antisense primer 3'AOXI (above). For the expression constructs containing a Myc tagged *Trichoderma reesei*  $\alpha$ -1,2-mannosidase expression unit, further evidence for genomic integration was obtained using Southern Blotting with the entire MFManMycHDEL ORF (<sup>32</sup>P labelled using HighPrime, Boehringer Mannheim) as a probe.

### 20 1.3 Expression of $\alpha$ -1,2-mannosidase

Expression of an  $\alpha$ -1,2-Mannosidase in GS115 strains expressing influenza virus haemagglutinin was verified by qualitative Northern blot. Expression of an  $\alpha$ -1,2-Mannosidase in PPY120H strains was verified by anti-Myc Western blot.

*Qualitative Northern Blot* -- Total RNA was purified from *Pichia* strains and the yield was determined spectrophotometrically. Northern blotting was performed according to standard procedures and an estimate of the quantity of RNA loaded was made using methylene blue staining of the blot, visualizing the rRNA

bands. The blot was probed with a *Clal/NarI* fragment of the mannosidase, labelled with  $^{32}\text{P}$  using HighPrime (Boehringer Mannheim).

*SDS-PAGE and Western Blotting* -- Total yeast cell lysates were prepared by washing the cells twice with PBS, followed by boiling in 1 volume of 2x concentrated Laemmli loading buffer for 5 min. The lysate was spun briefly in a microcentrifuge prior to gel loading and only the supernatant was loaded. For the analysis of proteins secreted into the growth media, the proteins were precipitated from 200  $\mu\text{l}$  of these media using desoxycholate/trichloroacetic acid according to standard procedures. The pellet was redissolved in 2x concentrated Laemmli loading buffer and the solutions were pH-corrected using Tris. SDS-PAGE was performed and followed by semidry electroblotting to nitrocellulose membranes. For Western Blotting, the 9E10 anti-Myc and the anti-HA mouse monoclonals (Boehringer Mannheim) were used at a concentration of  $1\mu\text{g/ml}$ , and the rabbit anti-PDI antiserum (Stressgen) was used at a dilution of 1/500. The secondary antibodies were goat anti-mouse IgG conjugated to alkaline phosphatase for the monoclonals and goat anti-rabbit IgG conjugated to peroxidase for the polyclonal (secondary antibodies from Sigma). Detection was performed using the NBT/BCIP system for alkaline phosphatase and the Renaissance substrate (NENBiosciences) for peroxidase. Imaging of the latter blot result was done on a Lumilager imaging device (Boehringer Mannheim).

The results shown in **Figure 4** indicated that the great majority of the HDEL-tagged protein was retained intracellularly, both when expressed from the strong constitutive GAP promoter and when expressed from the strong inducible AOXI promoter.

#### **1.4 Localization of $\alpha$ -1,2-Mannosidase**

*Isopycnic sucrose density gradient centrifugation* -- To determine the localization of the HDEL-tagged mannosidase, subcellular fractionation was carried

out using cells expressing the mannosidase-Myc-HDEL from the strong constitutive GAP promoter.

Briefly, 0.5 g of wet weight yeast cells were lysed using 4 x 1 min vortexing with 4.5 g glass beads in 1 ml lysis-buffer (50 mM Tris-HCL pH 7.5 containing 0.6 M sorbitol, 10 mM  $\beta$ -mercaptoethanol and 5 mM  $MgCl_2$ ). Between vortexing periods, the mixture was placed on ice for 5 min. The supernatant was collected and the glass beads were washed once with lysis-buffer, and the supernatant of this washing step was added to the first supernatant. This lysate was subjected to a differential centrifugation procedure. The P10000 pellet was solubilized in 0.5 ml of a 60% sucrose solution in lysis buffer. This solution was placed at the bottom of an Ultraclear ultracentrifuge tube (Beckman) of 14 x 89 mm. Subsequently, 1.5 ml each of sucrose solutions of 55, 50, 45, 42.5, 40, and 37.5% were carefully layered over each other. The tube was filled to the edge with 35% sucrose. Isopycnic sucrose gradient centrifugation was performed for 14 h at 180,000g in a Beckman SW 41 rotor in a Beckman Model L8-70 preparative ultracentrifuge. After completion, 1ml fractions were collected from the top and partially dialysed from excess sucrose, evaporated to dryness in a vacuum centrifuge. After redissolving the pellet in Laemmli buffer, the samples were subjected to SDS-PAGE in triplicate and the Western blots were treated with anti-HA, anti-Myc or anti-PDI ("PDI" for Protein Disulfide Isomerase), respectively.

The results illustrated almost exact cosedimentation of the MFManMycHDEL protein with the Protein Disulfide Isomerase marker protein (which is also targeted with a HDEL signal sequence) (**Figure 5**). In the same assay, the HA-tagged OCH1 was distributed over the whole gradient, with the highest abundance in fractions having a density lower than that of the fractions containing the mannosidase and the PDI. This result indicated that the mannosidase was targeted to the expected location (the ER-Golgi boundary) by the addition of an HDEL signal. In contrast, the mannosidase without HDEL, expressed from inducible alcohol oxidase I

promoter (which was of comparable strength as the GAP promoter), was secreted at a high level of about 20 mg/l.

*Immunofluorescence microscopy* -- To confirm the correct targeting of the mannosidase-Myc-HDEL, an immunofluorescence microscopy experiment was

5 performed.

Briefly, yeast cultures were grown to OD<sub>600</sub> in YPD (for pGAPZMFMManMycHDEL) or in YMP following a YPGlycerol growth phase for pPICZBMFMManMycHDEL. Formaldehyde was added to the yeast cultures to a final concentration of 4% and incubated for 10 min at room temperature. Cells were  
10 pelleted and resuspended in 50mM potassium phosphate buffer pH 6.5 containing 1mM MgCl<sub>2</sub> and 4% formaldehyde and incubated for 2h at room temperature. After pelleting, the cells were resuspended to an OD<sub>600</sub>=10 in 100 mM potassium phosphate buffer pH 7.5 containing 1mM MgCl<sub>2</sub> and EDTA-free Complete<sup>TM</sup> protease inhibitor cocktail (Boehringer Mannheim). To 100 µl of cell suspension, 0.6 µl of β-mercapto-  
15 ethanol and 20µl of 20,000 U/ml Zymolyase 100T (ICN) were added, followed by a 25 minute incubation with gentle shaking. The cells were washed twice in the incubation buffer and added to poly-lysine coated cover slips (these are prepared using adhesive rings normally in use for reinforcing perforations in paper). Excess liquid was blotted with a cotton swab and the cells were allowed to dry at 20°C. All  
20 blocking, antibody incubation and washing steps are performed in PBS containing 0.05% bovine serum albumin. Primary antibodies are used at 2µg/µl and secondary antibodies conjugated to fluorphores (Molecular probes) were used at 5µg/µl. The nucleus was stained with the nucleic acid stain HOECHST 33258. After fixation and cell wall permeabilization, the integrity of the yeast cell morphology was checked in  
25 phase contrast microscopy and after immunostaining, the slides were examined under a Zeiss Axiophot fluroresensce microscope equipped with a Kodak digital camera. Images were processed using Macprobe 4.0 software and prepared with Corel Photopaint 9.0.

The Golgi marker protein OCH1-HA gave the typical Golgi staining pattern described in the literature (speckle-like staining). Staining with the 9E10 monoclonal anti-Myc antibody, recognizing mannosidase-Myc-HDEL, gave a perinuclear staining pattern with some disparate staining in the cytoplasm, highly indicative for an ER targeting (Figure 4).

Based on the foregoing experiments, it is concluded that the *Trichoderma reesei* mannosidase-Myc-HDEL was targeted to the ER-Golgi boundary.

### Example 2

#### **Co-expression of Mannosidase-HDEL with Recombinant Glycoproteins**

##### Co-expression of Mannosidase-HDEL with the *Trypanosoma cruzi trans*-Sialidase

The cloning of a *Trypanosoma cruzi trans*-sialidase gene coding for an active *trans*-sialidase member without the C-terminal repeat domain has been described by Laroy et al. (*Protein Expression and Purification* 20: 389, 2000) which is incorporated herein by reference. The sequence of this *Trypanosoma cruzi trans*-sialidase gene is available through NCBI Genbank under the Accession No. AJ276679. For expression in *P. pastoris*, the entire gene was cloned in pHILD2 (Invitrogen, San Diego, CA), creating pHILD2-TS. To allow better secretion, pPIC9-TS was created in which *trans*-sialidase was linked to the prepro secretion signal of the yeast  $\alpha$ -mating factor. Plasmids pPIC9-TSE and pCAGGS-prepro-TSE were created where the epitope E-tag was added to the C-terminal of the *trans*-sialidase to allow easy detection and purification. The construction of pHILD2-TS, pPIC9-TSE and pCAGGS-prepro-TSE has been described by Laroy et al. (2000), incorporated herein by reference. The vectors used in the construction were made available through <http://www.belspo.be/bccm/lmbp.htm#main> for pCAGGS (No. LMBP 2453), Invitrogen, San Diego, CA for pHILD2 and pPIC9, and Pharmacia Biotech for pCANTAB-5E.



Plasmid pPIC9-TSE was linearized with SstI and was transformed into *P. pastoris* GS115 (*his4*) strain by electroporation according to the manufacturer's instructions (Invitrogen). One of the transformants was further transformed with plasmid pGAPZMFManHDEL, establishing a strain co-expressing Mannosidase-HDEL and the *Trypanosoma cruzi* *trans*-sialidase.

Fermentation and protein purification was according to the procedures described by Laroy et al. (2000).

Purified *trans*-sialidase was subject to carbohydrate analysis according to Callewaert et al., *Glycobiology* 11, 4, 275-281, 2001. Briefly, the glycoproteins were bound to the PVDF membrane in the wells of a 96-well plate, reduced, alkylated and submitted to peptide-N-glycosidase F deglycosylation. The glycans were derivatised with 8-amino-1,3,6-pyrenetrisulfonic acid by reductive amination. Subsequently, the excess free label was removed using Sephadex G10-packed spin columns and the glycans were analysed by electrophoresis on a 36 cm sequencing gel on an ABI 377A DNA-sequencer and detected using the built-in argon laser. Digests with 3 mU/ml purified *T. reesei*  $\alpha$ -1,2-mannosidase (described by Maras et al., *J. Biotechnol.* 77, 255-63, 2000) were also performed in 20 mM sodium acetate pH=5.0. The glycans derived from 1  $\mu$ g of the purified recombinant glycoproteins were used as the substrate. 1U of the  $\alpha$ -1,2-mannosidase is defined as the amount of enzyme that releases 1  $\mu$ mol of mannose from baker's yeast mannan per minute at 37°C and pH=5.0.

As can be seen in **Figure 6**, panel B, the major N-glycan on *trans*-sialidase was Man<sub>8</sub>GlcNAc<sub>2</sub> (Compare with panel F, representing an analysis of the N-glycans of bovine RNaseB. The one but last peak in this profile is Man<sub>8</sub>GlcNAc<sub>2</sub>, the first peak is Man<sub>5</sub>GlcNAc<sub>2</sub>). *In vitro*, this glycan was digestible to Man<sub>5</sub>GlcNAc<sub>2</sub> with  $\alpha$ -1,2-mannosidase (**Figure 6**, panel C). In the N-glycan profile of the *trans*-sialidase co-expressed with mannosidase-HDEL, the major peak corresponded to Man<sub>5</sub>GlcNAc<sub>2</sub> (**Figure 6**, panel D).

Co-expression of Mannosidase-HDEL with the Influenza A virus haemagglutinin

The Influenza A virus haemagglutinin was known to be glycosylated in *Pichia pastoris* with high-mannose N-glycans containing 9-12 mannose residues (Saelens et al. *Eur. J. Biochem.* 260: 166-175, 1999). The effect of a co-expressed  
5 mannosidase on the N-glycans of the haemagglutinin was assessed in an N-glycan profiling method described below. In addition, to compare the efficiency of the *Trichoderma* enzyme (having a temperature optimum of 60°C) with a mammalian mannosidase having a temperature optimum of 37°C, the catalytic domain of the mouse mannosidase IB from a mouse cDNA-library was cloned and tagged with a  
10 HDEL signal by PCR amplification. This ORF was cloned after the prepro-signal sequence of the *S. cerevisiae*  $\alpha$ -mating factor under the control of the GAP promoter. Expression of the mannosidase-HDEL transgenes on the mRNA level was confirmed by qualitative Northern blotting.

The haemagglutinin was expressed and purified from a non-mannosidase  
15 expressing control strain and from a strains co-expressing the *Trichoderma reesei* mannosidase-HDEL or the mouse mannosidase IB-HDEL according to the procedure described by Kulakosky et al. *Glycobiology* 8: 741-745 (1998). The purified haemagglutinin was subjected to PNGase F digestion as described by Saelens et al. *Eur. J. Biochem.* 260: 166-175, 1999. The proteins and glycans were precipitated with 3  
20 volumes of ice-cold acetone and the glycans were extracted from the pellet with 60% methanol. Following vacuum evaporation, the glycans were labeled with 8-amino-1,3,6 pyrenetrisulfonic acid by adding 1  $\mu$ l of a 1:1 mixture of 20 mM APTS in 1.2M citric acid and 1M  $\text{NaCNBH}_3$  in DMSO and incubating for 16h at 37°C at the bottom of a 250  $\mu$ l PCR-tube. The reaction was stopped by the addition of 10  $\mu$ l deionized  
25 water and the mixture was loaded on a 1.2 cm Sephadex G10 bed packed to dryness in a microspin-column by centrifugation in a swinging bucket rotor, which provided for a flat resin surface. After loading, 50  $\mu$ l deionised water was carefully added to the resin bed and the spin column was briefly centrifuged for 5 seconds at 750g in a tabletop centrifuge. This elution process was repeated twice and all the eluates were

pooled and evaporated to dryness in a Speedvac vacuum centrifuge (Savant). The labeled glycans were reconstituted in 1.5  $\mu$ l gel loading buffer containing 50% formamide and 0.5  $\mu$ l Genescan 500<sup>TM</sup>, labeled with rhodamine (Perkin Elmer Bioscience), serving as an internal reference standard. This mixture was loaded on a  
5 DNA-sequencing gel containing 10% of a 19:1 mixture of acrylamide:bisacrylamide (Biorad, Hercules, CA, USA) and made up in the standard DNA-sequencing buffer (89 mM Tris, 89 mM borate, 2.2 mM EDTA). Polymerization of the gel was catalyzed by the addition of 200  $\mu$ l 10% ammoniumpersulfate solution in water and 20  $\mu$ l TEMED. The gel was of the standard 36 cm well-to-read length and was run on  
10 an Applied Biosystems Model 373A DNA-sequencing apparatus. Prerunning of the gel was done at 1000 V for 15 min. and after loading, the gel was electrophoresed for 8h at 1250 V without heating. This methodology gives a limit of detection of 10 fmol per peak. The data were analysed with Genescan 3.0 software.

As shown in **Figure 7**, the *Trichoderma reesei*  $\alpha$ -1,2-mannosidase  
15 provided the most complete reduction in the number of  $\alpha$ -1,2-mannoses present on the N-glycans. The N-glycan processing by mouse mannosidase IB-HDEL was less efficient than by the *Trichoderma reesei*  $\alpha$ -1,2-mannosidase.

Despite the efficient removal of  $\alpha$ -1,2-mannoses from the N-glycans of haemagglutinin, no Man<sub>5</sub>GlcNAc<sub>2</sub> was obtained. Even after digestion of the N-  
20 glycans with 3 mU of purified *Trichoderma reesei*  $\alpha$ -1,2-mannosidase, only Man<sub>6</sub>GlcNAc<sub>2</sub> was obtained as the smallest sugar chain. These results indicated that the remaining residues were possibly  $\alpha$ -1,6-linked mannoses, originating from the initiating OCH1  $\alpha$ -1,6-mannosyltransferase enzymatic activities. OCH1 was observed to be localized to very early part of the Golgi apparatus and could act on the N-glycans  
25 of haemagglutinin before complete digestion of the Man<sub>8</sub>GlcNAc<sub>2</sub> precursor to Man<sub>5</sub>GlcNAc<sub>2</sub> by the mannosidases-HDEL. Thus, for proteins whose glycans are efficiently modified by the  $\alpha$ -1,6-mannosyltransferase, an inactivation of the OCH1

gene coding for the transferase would be desirable in order to obtain proteins with  $\text{Man}_5\text{GlcNAc}_2$ .

### Example 3

#### **Inactivation of the *Pichia* Och1 Gene:**

A *Pichia pastoris* sequence was found in the GenBank under Accession  
5 No. E12456 and was described in Japanese Patent Application No. 07145005,  
incorporated herein by reference. This sequence shows all typical features of an  $\alpha$ -  
1,6-mannosyltransferase and is most homologous to the *S. cerevisiae* OCH1, thus  
referred to herein as the *Pichia pastoris* Och1 gene.

First, the full ORF of the *Pichia pastoris* Och1 gene was PCR cloned in  
10 pUC18 to obtain plasmid pUC18pOch1. pUC18pOch1 was cut with HindIII, blunt-  
ended with T4 polymerase, then cut with XbaI, releasing a fragment containing the 5'  
part of the *Pichia pastoris* Och1 gene. This fragment was ligated into the vector  
pBLURA IX (available from the Keck Graduate Institute, Dr. James Cregg,  
<http://www.kgi.edu/html/noncore/faculty/cregg/cregg.htm>), which had been cut with  
15 *Eco* RI, blunt-ended with T4 polymerase, and then cut with *Nhe* I. This ligation  
generated pBLURA5'PpPCH1, as shown in **Figure 8**.

Disruption of this *Pichia* OCH1 gene in the *Pichia* genome was achieved  
by single homologous recombination using pBLURA5'PpOCH1, as illustrated in  
**Figure 9**. As a result of the single homologous recombination, the Och1 gene on the  
20 *Pichia* chromosome was replaced with two Och1 sequences: one consisted only about  
the first one third of the full Och1 ORF, the other had a full Och1 ORF without a Och1  
promoter. Single homologous recombination was achieved as follows. Cells of the  
*Pichia* strain yGC4 were transformed by electroporation with pBLURA5'PpOCH1  
which had been linearized with the single cutter *Bst* BI. About 500 transformants  
25 were obtained on minimal medium containing 1M sorbitol, biotin, arginine, adenine  
and histidine and incubation at 27°C. Thirty-two of these transformants were picked  
and re-selected under the same conditions. Twelve clones were further analyzed for  
correct genomic integration of the cassette by PCR. Seven of the twelve URA  
prototrophic clones contained the cassette in the correct locus.

One of the Och1-inactivated clones was also further transformed with pGAPZMFManHDEL to produce “supertransformants”. Both the Och1-inactivated clone and three supertransformants also expressing the ManHDEL were evaluated in cell wall glycan analysis as follows. Yeast cells were grown in 10 ml YPD to an  
5 OD<sub>600</sub>=2 and mannoproteins were prepared by autoclaving the yeast cells in 20 mM sodium citrate buffer pH7 for 90 min at 120 °C and recovery of the supernatant after centrifugation. Proteins were precipitated from this supernatant with 3 volumes of cold methanol. The protein preparation obtained in this way was used for N-glycan analysis using DSA-FACE as described by Callewaert et al. (2001) *Glycobiology* 11,  
10 275-281. As shown in **Figure 10**, there was an increased amount of Man<sub>8</sub>GlcNAc<sub>2</sub> glycan in the Och1-inactivated clone as compared to parent strain yGC4, indicative of a reduced activity of the Och1 enzyme. In all three supertransformants which also expressed the HDEL-tagged  $\alpha$ -1,2 mannosidase, the production of Man<sub>5</sub>GlcNAc<sub>2</sub> was observed. Furthermore, upon digestion of the same glycan mixtures with 3 mU/ml  
15 purified recombinant *Trichoderma reesei*  $\alpha$ -1,2-mannosidase, more Man<sub>5</sub>GlcNAc<sub>2</sub> was formed in the strain transformed with pBLURA5'PpOCH1 than in the parent strain (**Figure 11**, compare panel 2 and 3).

These results confirmed that the lack of a production of Man<sub>5</sub> glycans on recombinantly produced proteins such as haemagglutinin from cells expressing  $\alpha$ -1,2-  
20 mannosidase were due to the activity of the Och1 protein. These results further indicate that the production of glycoproteins with Man<sub>5</sub> glycans could be facilitated by the inactivation of the Och1 gene.

### Example 4

### Expression of Glucosidase II in *Pichia pastoris*

#### **4.1 Amplification of the GLSII alpha subunit ORF from *S. cerevisiae*.**

5 Genomic DNA was prepared from the *S. cerevisiae* strain INVS ( $\alpha$ , leu2-3, 112 his3 $\Delta$ 1, trp1-289, ura3-52), using the Nucleon kit (Amersham). A touch-down PCR reaction was performed using this genomic DNA as template and the LA TaKaRa polymerase (ImTec Diagnostics). The sequence of the PCR primers was based on the known sequence of the *S. cerevisiae* GLSII ORF:

10 Sense primer: 5' CCG CTC GAG ATG GTC CTT TTG AAA TGG CTC 3'

(SEQ ID NO:12)

Antisense primer: 5' CCG GGC CCA AAA ATA ACT TCC CAA TCT TCA  
Apa I

15 G 3' (SEQ ID NO:13)

## **4.2 Cloning of the *S. cerevisiae* glucosidase II ORF into *Pichia pastoris* expression vectors.**

*Construction of the glucosidase II expression vectors* -- The PCR fragment was digested with *Xho* I/*Apa* I and ligated into the pGAPZA vector (Invitrogen), thereby placing the ORF under the transcriptional control of the GAP promoter. Using this strategy, the myc and the His6 tag were placed in frame to the C-terminus of Glucosidase II, creating pGAPZAGLSII. The complete ORF of pGAPZAGLSII was then sequenced to ensure that no mutations were generated in the PCR reaction.

The sequence of the vector pGAPZAGLSII was set forth in SEQ ID NO: 18. The GLSII ORF from the pGAPZAGLSII vector was cloned into vector pPICZA (Invitrogen) to create pPICZAGLSII, thereby placing the ORF under the transcriptional control of the AOXI promoter. The GLSII ORF from the pGAPZAGLSII vector was cloned into vector pAOX2ZA, thereby placing the ORF under the transcriptional control of the AOX2 promoter. This vector was created by

replacing the multi cloning site of vector pAOX2ZB with the multi cloning site of pPICZA. Vector pAOX2ZB was generated by replacing the AOX1 promoter of pPICZB by the AOX2 promoter region of the AOX2 gene (Martinet et al., Biotechnology Letters 21). The AOX2 promoter region was generated by PCR on

5 *Pichia* genomic DNA with the sense primer  
 5'GACGAGATCTTTTTTTCAGACCATATGACCGG 3' (SEQ ID NO: 26) and the  
 antisense primer 5'GCGGAATTCTTTTCTCAGTTGATTTGTTTGT 3' (SEQ ID  
 NO: 27). The GLSII ORF from the pGAPZGLSII vector was cloned into vector  
 pYPT1ZA to create pYPT1ZAGLSII, thereby placing the ORF under the

10 transcriptional control of the YPT1 promoter. Vector pYPTZA was created by  
 replacing the AOX1 promoter of pPICZA by the YPT1 promoter present on the  
 plasmid pIB3 (GenBank accession number AF027960)(Sears et al., Yeast 14, pg 783-  
 790, 1998). All constructs contain the phleomycin resistance gene. The resulting final  
 expression vectors (pGAPZAGLSII, pAOX2ZAGLSII, pPICZAGLSII and

15 pYPT1ZAGLSII) are depicted in **Figures 12-15**.

Similar expression vectors were constructed, carrying the Ampicillin  
 resistance marker and the *Pichia* ADE1 selection marker. In principle, the Zeocin  
 resistance expression cassette of the plasmids pAOX2ZAGLSII, pGAPZAGLSII and  
 pYPT1ZAGLSII was replaced by the Ampicillin and *Pichia* ADE1 cassette of the

20 vector pBLADE IX (Cregg, J.M.) to result in the vectors pAOX2ADE1glsII,  
 pGAPADE1glsII and pYPT1ADE1glsII. Vector pPICADE1glsII was obtained by  
 inserting the glucosidase II open reading frame into the multiple cloning site of the  
 vector pBLADE IX (Cregg, J.M.). The resulting final expression vectors  
 (pGAPADE1glsII, pAOX2ADE1glsII, pPICADE1glsII and pYPT1ADE1glsII) are

25 depicted in **Figures 16-20**.

*Adding the ER retention tag HDEL to Glucosidase II expression vectors –*

The following primers were used to generate an HDEL-containing PCR fragment:

Primer 1: 5'GCG GGT CGA C/CA C/GA C/GA A/CT G/TG A/GT TTT AGC CTT  
                                     Sal I       H       D       E       L       stop



AGA CAT GAC 3' (SEQ ID NO:28)  
Primer 2: 5'CAG GAG CAAA GCT CGT ACG AG 3' (SEQ ID NO:29)  
*Spl* I

5 PCR was performed on pGAPZAGLSII with Taq pol., at 60°C. The PCR  
fragment of 225 bp was cut with *Sal* I/*Spl* I and ligated into the *Sal* I/*Spl* I opened  
pGAPZAGLSII vector, creating plasmid pGAPZAglslIIHDEL. The sequence of  
plasmid pGAPZAglslIIHDEL is set forth in SEQ ID NO: 24. The construction  
strategy and the resulting final expression vectors (pGAPZAglslIIHDEL and  
10 pGAPADE1glslIIHDEL) are depicted in **Figures 20-21**.

### 4.3 Transformation of a *Pichia pastoris* strain.

Transformation was performed using the conventional electroporation techniques, as described by Invitrogen. Cells of the *Pichia pastoris* strain PPY12-OH  
15 were transformed with pGAPZGLSII which had been cut with the single cutter *Avr* II. Transformants were selected based on their resistance to zeocin.

*Genomic analysis of the transformants* -- Genomic DNA was prepared from some zeocin resistant *Pichia* transformants. A PCR reaction was performed on the genomic DNA in order to determine whether or not the glucosidase II gene was integrated into the yeast genome. PCR was performed using Taq DNA polymerase (Boehinger) (2.5 mM MgCl<sub>2</sub>, 55°C for annealing). The primers were the same as the ones we used for the amplification of the ORF on *S. cerevisiae* genomic DNA. pGAPZAGLSII transformants were confirmed by the presence of a specific PCR product indicative of the glucosidase II ORF.

#### **4.4 Expression and secretion of the *S. cerevisiae* glucosidase II alpha subunit in *Pichia pastoris***

*Analysis at the transcriptional level* -- RNA was prepared from the transformants which scored positive after the genomic analysis. RNA was prepared using acid phenol. From each sample, 15 µg of RNA was loaded on a formaldehyde

agarose gel. After electrophoresis the RNA was blotted on a Hybond N membrane. The membrane was hybridizing using a radioactive probe, which consists of a 344 bp glucosidase II specific fragment, corresponding to the 3' region of the glucosidase II ORF. No signals were detected with non-transformed control strains, whereas clear  
5 signals were observed with transformants.

*Analysis at the protein level using a double membrane assay -- A*

nitrocellulose membrane was placed on a buffered dextrose medium (BMDY). On top of that nitrocellulose membrane, a cellulose acetate membrane was placed. *Pichia* transformants of pGAPZAGLSII were streaked on the cellulose acetate and grown for  
10 a few days. The yeast cells remained on the cellulose acetate, while the secreted proteins crossed this membrane. As such the secreted protein was captured onto the nitrocellulose membrane. After a few days the cellulose acetate, containing the yeast colonies, was removed. The nitrocellulose membrane was analyzed for the presence of glucosidase II using anti-myc antibody. Most of the transformants gave a clear  
15 signal as compared to a faint, hardly visible signal with the WT, non-transformed strain.

*Extracellular expression* – PPY12-OH transformants of the construct pGAPZAGLSII(mychis6) (strains 12, 14 and 18) and transformants of the construct pGAPZAGLSII(myc)HDEL (strains H1, H2 and H3) were grown for 2 days on 2x10  
20 ml BMDY medium. These 6 transformants earlier scored positive both on the genomic level (PCR on gDNA) and on the RNA level (Northern blot). The culture medium was collected by centrifugation and concentrated with Vivaspin columns to about 1 ml. Proteins from this concentrate were precipitated with TCA, resuspended in Laemmli buffer and loaded for SDS-PAGE analysis. Proteins were blotted to  
25 nitrocellulose membrane. The blot was incubated overnight with anti-myc Ab. The secondary Ab was linked to peroxidase. Using the Renaissance luminiscence detection kit (NEN) and a light sensitive film (Kodak), a strong band at about 110 kDa was observed for the transformants 12, 14 and 18, indicating that GLSII was expressed and secreted from these transformants. No signal was obtained for the transformants

H1-3, which indicate that the HDEL tag, which was added C-terminally to the GLSII ORF, resulted in an ER localization of the protein, preventing GLSII to be secreted into the growth medium.

*Intracellular expression* – The 6 transformants and the WT strain were grown for 2 days in 500 ml BMDY. The cells were collected by centrifugation, washed, resuspended into a minimal volume (50 mM Tris.HCl pH 7.5, 5% glycerol) and broken using glass beads. The cell debris was removed through several centrifugation steps (low speed centrifugation (2000-3000g)). Membranes were obtained from the supernatant through ultracentrifugation. The pellets were resuspended in Laemmli buffer and loaded for SDS-PAGE analysis. The proteins were blotted on a nitrocellulose membrane. The intracellular GLSII expression was checked using anti-myc Ab and peroxidase conjugated secondary Ab. Following the luminescence detection, a band at about 110 kDA was observed with the GLSIIHDEL tranformants (H1 and H3, faint signal for H2), but not with the WT and GLSII expression strains. These results clearly indicate the intracellular presence of the recombinant GLSII when expressed with a C-terminal HDEL tag. No GLSII was detected intracellularly when this tag was not present.

#### **4.5 Purification and activity assays of the recombinant glucosidase II alpha**

**submit**

A GLSII assay was set up as follows and was tested using a commercially available yeast alpha-glucosidase (Sigma) as a positive control.

Composition: 70 µl 80 mM phosphate-citrate buffer pH 6.8, 7 µl 250 mM mannose, 3.5 µl 250 mM 2-deoxy-D-glucose, 0.8 µl 4-MeUmbelliferyl-alpha-D-glucopyranoside (1 µM). Three assays were performed: one with 1 unit commercial enzyme, one without the enzyme and one with the enzyme but without the substrate. The assay mixture was incubated overnight at 30°C. When illuminated with UV, only the reaction mixture with both the enzyme and the substrate showed fluorescence

(Figure 22). This indicates that the assay was very specific in detecting the activity of the alpha-glucosidase.

- WT PPY12-OH, strain 18 and strain H3 were grown during 2 days in 2x10 ml growth medium. Cells were spun down and medium was adjusted to 300 mM NaCl and 10 mM imidazol and concentrated with Vivaspin columns to 0.5-1ml. Medium was loaded onto a Ni-NTA spin column (Qiagen) and the purification was performed according to the manufactures recommendations. Protein was eluted from the column in 2x100 µl elution buffer (50 mM NaH<sub>2</sub>PO<sub>4</sub>, 300 mM NaCl, 250 mM imidazol pH 8.0). From each eluate, 20 µl was assayed for its glucosidase II activity.
- 0.33 units of the commercial enzyme diluted in 20 µl of the elution buffer was used as a positive control. Fluorescence was observed with the positive control and the elute of strain 18, the strain which secreted the enzyme into the growth medium. These results indicate that the recombinant *S. cerevisiae* GLSII alpha subunit, secreted by *Pichia pastoris*, was a functionally active enzyme. The activity was not seen in the WT (untransformed) strain, nor in strain H3 as the GLSII was expressed intracellularly (Figure 23). These results also indicate that the beta subunit is not necessary for the functionality of the alpha part of the protein.

**Example 5**  
**Creating *Pichia* Strains Expressing  
both Glucosidase II and Mannosidase**

5                Strain GS115 was transformed with pGAPZGLSII and pGAPZglsIIHDEL.  
Transformants were selected on YPDSzeo.

              Strain yGC4 was transformed with the following constructs, respectively:

- (1) pGAPADEglsII and pGAPADEglsIIHDEL, selection on synthetic  
sorbitol medium without adenine;
- 10            (2) pGAPZMFManHDEL: selection on YPDSzeo; and
- (3) pGAPZMFManHDEL/pGAPADEglsIIHDEL: selection on synthetic  
sorbitol medium without adenine and with zeocin.

              Strain yGC4 with OCH1 knock-in and expressing MFmannosidaseHDEL  
was transformed with pGAPADEglsII and pGAPADEglsIIHDEL. Selection of

15            transformants was done on synthetic sorbitol medium without adenine and uracil.

              For all transformations, colonies were obtained. Transformants with the  
expression vector(s) integrated into the genome, determined by PCR, were obtained.  
Expression of GLSII from some of these transformants was observed.

20

## SEQUENCE LISTING

- SEQ ID NO: 1**  
HDEL (peptide)  
5
- SEQ ID NO: 2**  
5'-GACTGGTTCCAATTGACAAGC-3'
- SEQ ID NO: 3**  
5'-AGTCTAGATTACAACCTCGTCGTGAGCAAGGTGGCCGCCCCG TCG-3'
- 10 **SEQ ID NO: 4**  
CCATTGAGGACGCATGCCGCGCC
- SEQ ID NO: 5**  
GTATCTAGATTACAACCTCGTCGTGCAGATCCTCTTCTGAGATGAGTTTTTGT  
15 TCAGCAAGGTGGCCGCCCCGTCGTGATGATGAA
- SEQ ID NO: 6**  
AACTCGAGATGGACTCTTCAAAACACAAACGC
- 20 **SEQ ID NO: 7**  
TTGCGGCCGCTTACAACCTCGTCGTGTCGGACAGCAGGATTACCTGA
- SEQ ID NO: 8**  
CCATTGAGGACGCATGCCGCGCC  
25
- SEQ ID NO: 9**  
GCAAATGGCATTCTGACATCCT
- SEQ ID NO: 10**  
30 GTCCCTATTTCAATCAATTGAA
- SEQ ID NO: 11**  
GACTGGTTCCAATTGACAAGC
- 35 **SEQ ID NO: 12**  
CCG CTC GAG ATG GTC CTT TTG AAA TGG CTC
- SEQ ID NO: 13**  
40 CCG GGC CCA AAA ATA ACT TCC CAA TCT TCAG

**SEQ ID NO: 14**

The ORF sequence of the MFManHDEL fusion in pGAPZMFManHDEL:

atgagatttccttcaattttactgctgtttattcgcagcatcctccgattagctgctccagtcaacactacaacagaagatgaa  
 5 acggcacaaattccggctgaagctgtcatcggttactcagatttagaaggggatttcgatgttgctgtttgccattttcaaca  
 gcacaaataacgggtattgtttataataactactattgccagcattgctgctaaagaagaagggtatctctcgagaaaagag  
 aggctgaagctgaattcgccacaaaacgtggatctccaaccctacgagggcggcagcagtcaggccgcatccagac  
 gtcgtggaacgcttaccaccattttgcctttcccatgacgacctccaccggtcagcaacagcttgatgatgagagaaacg  
 gctggggctcgtcggcaatcgatggcttggacacggctatcctcatgggggatgccgacattgtgaacacgacacctcagta  
 10 tgtaccgcagatcaactcaccacgactgcggttccaaccaaggatcctccgtgttcgagaccaacattcggtacctcgggt  
 ggctgctttctgcctatgacctgttgcgaggtccttcagctccttggcgacaaaccagaccctgtaaacagccttctgag  
 gcaggctcaaacactggccaacggcctcaaggttgcgttcaccactcccagcgggtgtcccggaccctaccgttcttcaac  
 cctactgtccggagaagtgggtgcatctagcaacaacgtcgtgaaattggaagcctgggtgctcagtggtgacacgggtgagc  
 gacctgacgggaaaccgcagtatgccagcttgcgcagaagggcgagtcgtatctcctgaatccaaagggaagcccg  
 15 aggcattggcctggcctgattggaacgtttgcagcacgagcaacggctaccttcaggatagcagcggcagctggctccggc  
 ctcatggacagcttctacgagtacgtatcaagatgtacctgtacgacctgggtgcgtttgcacactacaaggatcgctgggt  
 ccttgggtccgactcgaccttgggcatctcggctctacccgtcgacgcgcaaggacttgaccttttgtcttcgtacaacg  
 gacagtctacgtcgccaaactcaggacatttggccagtttggcggtggcaacttcatttgggaggcatttctcctgaacgag  
 caaaagtacattgactttggaatcaagcttgcagctcgtactttggcacgtacaccagacggcttctggaatcgggcccgga  
 20 aggcttcgcgtgggtggacagcgtgacggggcgccggcggtcgcgccctcgtccagtcggggttctactcgtcggca  
 ggattctgggtgacggcaccgtattacatcctgcggccggagacgctggagagcttgactacgcataccgcgtcacggg  
 cgactccaagtggcaggacctggcgtgggaagcgttgagtccattgaggacgcatgccgcgccggcagcgcgtactcg  
 tccatcaacgacgtgacgcaggccaacggcggggggtgcctctgacgatatggagagcttctggtttgccgaggcgtcaa  
 gtatgcgtacctgatctttgcggaggagtcggatgtgcaggtgcaggccaccggcggaacaaattgtctttaacacgga  
 25 ggcgaccccttagcatccgttcatcatcacgacggggcgccaccttgctcacgacgagttgaa

**SEQ ID NO: 15**

The ORF sequence of the MFmManHDEL fusion in pGAPZMFmManHDEL:

30 atgagatttccttcaattttactgctgtttattcgcagcatcctccgattagctgctccagtcaacactacaacagaagatgaa  
 acggcacaaattccggctgaagctgtcatcggttactcagatttagaaggggatttcgatgttgctgtttgccattttcaaca

gcacaaataacgggttattgtttataaatactactattgccagcattgctgctaagaagaaggggtatctctcgagatggact  
 ctcaaaacacaaacgcttgatctgggcttagaagatgtgttaattcctcacgtagatccgggcaaaggagctaaaaacccc  
 ggcgcttctgatccatggacccgacgaacacagacacaggggaagaagaagagcgctcgagaaataagattagagctg  
 accatgagaaagccctggaagaagcaaaagaaaaattaagaaagtcaagagaggaaatccgtgcagaaattcagacaga  
 5 gaaaaacaaagtagcccaagcaatgaagacaaaagagaccagggtactgccgcctgtccctgtcccacaacgtgtaggg  
 gtcagtgggtggggatccagaagacatggagatcaagaagaaaagagacaaaattaagagatgatgaacatgcctggg  
 ataattacagaacatacggatggggacataatgaactaaggcctattgcaaggaaaggccattccactaacatattcggaag  
 ctcacagatgggtgccaccatagtggtatgcttggataccctttatatcatggggcttcatgatgaattcatggatgggcaaag  
 atggattgaagaaaaccttgattcagtgtgaattcagaagtgtctgtcttgaagttaacattcgtttattggagggctcctcg  
 10 ctgcatattacctgtcaggagaggaatatcaagactaaagcagtgagtggtgagaaactccttctgcctttaacacac  
 ctactgggattccctgggcaatggtgaacctgaaaagtgagtaggtcgaaactggggctggcgctgcaggcagcagc  
 atcctggctgagttcggcaccctgcacatggagttgtgcacctcagctacttgaccggtgacttgacttactataataaggtc  
 atgcacattcggaaactactgcagaaaatggaacgccc aaatggtctttatccaaattatttaaacccaagaacagggcgctg  
 gggtcagtatcacacatcagttggtggtctgggagatagttttatgaatacttactgaaagcatggctgacgtcagataaaac  
 15 agaccacgaggcaagaaggatgtatgacgatgctgtgaggtatagaaaaacatcttattaagaagtcccgaggaggtct  
 ggtttttattggagaatggaagaatggacacttggaaggaagatggggcacttgccctgcttctgctgggggaatgcttgcc  
 cttggagcagatggttcagaaaggataaagctggccactacttagaactaggggcagaaattgcacgaacatgtcatgag  
 tcatatgacagaactgcattgaaactaggtccggagtcattcaagttgatggtgcagtggagccgtggctgtgcggcagg  
 ctgaaaagtattacatccttcgtccagaagtaattgaaacctattggtatctatggcgatttaccacgaccaagatacaggc  
 20 agtggggctgggaagcagcactggctattgagaagtcgtccgggtcagcggtgggttttctggtgtcaaggatgtatacg  
 ccccgaccctgtgcatgacgacgtgcagcagagcttttcttctgctgaaacattaaaatacttgtacctgctgttctctggcga  
 tgaccttctacctttagaccactgggtgtttaacacagaggcgaccctctgccggtgttgcttagccaacagcactcttc  
 aggtaatcctgctgtccgacacgacgagttgtaa

25 **SEQ ID NO: 16** pAOX2ZAGLSII

catggccaagtgaccagtgccgttccggtgctcaccgcgcgcgacgtcggcgagcggtcgagttctggaccgaccgg  
 ctcgggttctccgggacttcgtggaggacgacttcgccggtgtggtccgggacgacgtgaccctgttcatcagcgcggtc  
 caggaccaggtggtgccggacaacacctggcctgggtgtgggtgcgcggcctggacgagctgtacgccgagtggtgcg  
 gaggtcgtgtccacgaactccgggacgcctccggccggccatgaccgagatcgccgagcagccgtgggggcggga



gttcgccctgcgcgacccggccggcaactgcgtgcacttcgtggccgaggagcaggactgacacgtccgacggcggcc  
cacgggtcccaggcctcggagatccgtcccccttttccttgcgatatcatgtaattagttatgtcacgcttacattcacgcct  
ccccccacatccgctctaaccgaaaaggaaggagtagacaacctgaagcttaggtccctatttttttatagttatgtagt  
attaagaacgttatttatattcaaattttctttttctgtacagacgcgtgtacgcatgtaacattatactgaaaaccttgcttga  
5 gaaggtttgggacgctcgaaggctttaattgcaagctggagaccaacatgtgagcaaaaggccagcaaaaggccagga  
accgtaaaaaggccgcttgcgtggcgttttccataggtccgccccctgacgagcatcacaaaaatcgacgctcaagtca  
gaggtggcgaaaccgacaggactataaagataccaggcgttccccctggaagctccctcgtgcgctcctgttccgac  
cctgccgcttaccggatacctgtccgcctttctcccttcgggaagcgtggcgctttctcaatgtcacgctgtaggtatctcagt  
tcggtgtaggtcgttcgctccaagctgggctgtgtgcacgaacccccgttcagcccgaccgctgcgccttatccgtaact  
10 atcgtcttgagtccaacccgtaagacacgacttatgccactggcagcagccactgtaacaggattagcagagcgaggt  
atgtaggcggtgctacagagtictgaagtgggtggcctaactacggctacactagaaggacagtatttggatctgcgctctg  
ctgaagccagttaccttcggaaaaagagttgtagctcttgatccggcaaacaaaccaccgctggtagcggtgtttttgtt  
tgcaagcagcagattacgctcagaaaaaaaggatctcaagaagatccttgatcttttctacggggctgacgctcagtggga  
acgaaaaactcacgttaagggttttggctatgagatcagatctttttcagaccatatgaccggctccatcttctacggggggat  
15 tatctatgctttgacctctatcttgattctttatgattcaaatcacttttacgttatttacttactggttatttacttagcgcttttctg  
aaaaacatttactaaaaatcatacatcggcactctcaaacacgacagattgtgatcaagaagcagagacaatcaccactaag  
gttgcacatttgagccagtaggctcctaatagaggttcgatacttattttgataatacgacatattgtcttacctctgaatgtgtca  
atactctctcgttctcgtcgtcagctaaaaatataacacttcgagtaagatacgcccaattgaaggctacgagataccaga  
ctatcactagtagaactttgacatctgctaaagcagatcaaatatccattatccagaatcaattaccttcccttagcttgcgaag  
20 gcatgaaaaagctacatgaaaatccccatccttgaagttttgcagcttaaaggactccatttctaaaaattcaagcagtcctct  
caactaaatttttccattcctctgcacccagccctctcatcaaccgtccagccttctcaaaagtccaatgtaagtagcctgca  
aattcaggttacaacccctcaattttccatccaaggggcgtaccttacaaagttaatatcgaacagcagagactaagcgagtcac  
catcaccaccaacgatgggtgaaaaacttaagcatagattgatggagggtgtatggcacttggcggctgcattagagtttga  
aactatggggtaatacatcacatccggaactgatccgactccgagatcatatgcaagcacgtgatgtacccgtaaaactgc  
25 tcggattatcgttgcaattcatcgtcttaaacagtacaagaaactttatcatgggtcattggactctgatgaggggcacatttcc  
ccaatgatttttgggaaagaaagccgtaagaggacaggttaagcgaagagacaagacaacgaacagcaaaagtacag  
ctgtcagctacctagtggacagttgggagtttccaattgggttggtttgaattttaccatgttgagttgtccttgccttctccttga  
aacaatgcaagttgataagacatcaccttcaagataggtatttttgcgcataaattttgtctcggagtgaaaacccctttat  
gtgaacagattacagaagcgtcctacccttcaccgggtgagatggggagaaaattaagcgtgaggagacgattattggtat

aaaagaagcaacaaaatcccttattgtcctttctgatcagcatcaaagaatattgtcttaaacgggctttaactacattgttc  
ttacacattgcaaacctctccttctatttcggatcaactgtattgactacattgatcttttaacgaagttacgacttactaaatcc  
ccacaaacaaatcaactgagaaaagaattcacgtggcccagccggccgtctcggatcggtagctcgcgatggctctttga  
aatggctcgtatgccaatgggtcttcttaccgcttttcgcatgcgtttaccgactatctattaagaagtgtgcgaatctgggt  
5 tttgccatagaaacagggttatgcagaaaatattgccaaatctcatcactgctattacaaagtggacgccgagtctattgcac  
acgatcctttagagaatgtgcttcatgctaccataaataaaactataccaagattggaggggcgatgatatagccgttcagttccc  
attctctctctctttttacaggatcactcagtaagggtcactataaatgagaaagagagaatgccaaccaacagcagcggtttg  
ttgatctcttcacaacgggtcaatgagacctggaagtacgcattcgacaagaaattcaaggaggggcgaacaggaccagta  
ttccacaattccacttccctaagcaaaaaacaaactgtgaactcattctggtcgaaaatatcttcattttgtcactttcaaacccac  
10 tgcagacacatttcatcttcgaaacgggtgatgtatccgtagaaatctttgctgaacctttcaattgaaagttactggcaaatg  
cgctgaaacttattgtaaacgagcaaaatttctgaacattgaacatcatagaactaagcaggaaaacttcgcacacgtgctg  
ccagaagaaacaactttcaacatgtttaaggacaatttctgtattcaagcatgactctatgcccttggggcctgaatcggttg  
cgctagatttctcttcatgggttctactaatgtctacgggtataccgggaacatgcgacgtcgctaaggctgatggacacttcagg  
tggaaaggaaccctacaggcttttaacgttgatgtctttgagtacaacatcggtaccagccaacaaatgtacgggttcgatcc  
15 caticatgtttcatcttcgtccacatctatctttgggtcaatgcagctgacacttgggtagacataaagtatgacaccagtaaa  
aataaaacgatgactcattggatctccgaaaatgggtgcatagatgtagtcatgtccctggggccagatatccaactatcattg  
acaaatttaccgattgactggtagacccttttaccgcccatttctctatagggtaccatcaatgtagatggaattataatgatg  
agatggacgttctcacagtggactctcagatggatgctcatatgattccttacgattttatttgggttgacttggagtatacgaac  
gacaaaaaatattttacttgaagcagcactccttccaatccaaaaaggctgttatccaaattaaaaaagggtgggtagaat  
20 ctgtcgtactaatcgatcctcatttaaagaaaagattatgaaatcagtgacagggttaattaatgaaatgtagcagtcaggatc  
acaatggaaatgactatgtaggctcattgctggccaggtaattctatatggattgataccataagcaaatatggccaaaagattt  
ggaagtccttttcgaacgggttatggatctgccggctgatttaactaatttattcatttggatgatatgaacgagccttcgatttt  
cgatggcccagagaccacagctccaaaagatttgattcacgacaattacattgaggaaagatccgtccataacatatatggtc  
tatcagtgcatgaagctacttacgacgcaataaaatcgattttaccatccgataagcgtccttcttctaacaagggttttt  
25 ttgccggctctcaacgtactgctgccacatggactggtgacaatgtggccaattgggattacttaagatttccattcctatggt  
tctgtcaaacaacattgctggatgccatttataggagccgacatagctggcttctgctgaggatcctacacctgaattgattgca  
cgttggtagcaagcgggcttatggtacccatttttagagcacacgcccatatagacaccaagagaagagaaccatacttatt  
caatgaacctttgaagtcgatagtacgtgatattatccaattgagatatttctgctacctaccttataccatgtttcataaatca  
agtgtcactggatttccgataatgaatccaatgtttattgaacacctgaatttctgaattgtatcatatcgataaccaattttact

ggagtaattcaggtctattagtcaaaccgtcacggagcctggtaacagaaacggaaatggtttcccacccggtatattct  
atgaattcgcacatctttacacatctttataaacaatggtactgatttgatagaaaagaatatttctgcaccattggataaaattccatt  
atttattgaaggcgggtcacattatcactatgaaagataagtagaagatcttcaatgttaatgaaaaacgatccatatgtaatag  
ttatagcccctgataccgaggagcagccgttgagatctttatgttgatgaggagaaacttttgctaccaaaagaggtgag  
5 tacgtagaaactcagttcattttcgaacaataaccttaaaaaatgttcgaagtcatttcccagaaatttgacaggcattcacc  
acaatactttgaggaataccaatattgaaaaatcattatcgcaagaataattacaacacataacgttgaaagacagta  
ttaaagtcaaaaaaaatggcgaagaaagttcattgccgactagatcgcatatgagaatgataataagatcaccattcttaacc  
tatcgcttgacataactgaagattgggaagttattttgggcccgaacaaaactcatctcagaagaggatctgaatagcgc  
gtcgaccatcatcatcatcattgagtttagccttagacatgactgttctcagttcaagttgggcacttacgagaagaccg  
10 gtcttgctagattctaataagaggatgtcagaatgccatttgccgtgagagatgcaggcttcattttgatactttttatttgaac  
ctatatagtataggattttttgtcattttgttcttctctgacgagcttgcctgatcagcctatctcgcagctgatgaatatctgt  
ggtaggggttggaataacattcgagttgatgttttcttggtatttccactcctctcagagtacagaagattaagttagac  
cttcgtttgtcggtatccccacacacatagctcaaaatgtttctactcctttttactcttccagattttctcgactccgcga  
tcgccgtaccacttcaaaacaccaagcacagcactaataatttccctcttcttctctaggggtgcgttaattaccgacta  
15 aagggttgaaaaagaaaaagagaccgcctgtttcttttctctgctgaaaaaggcaataaaattttatcacgtttcttttctt  
gaaatttttttttagtttttcttctcagtgacctcattgatattaaagtaataaacggcttcaatttctcaagtttcagtttcatt  
ttctgttctattacaacttttttacttctgttcattagaaagaaagcatagcaatctaagggcggtgtgacaattaatcat  
cggcatagtatcggcatagtataatacgaacaaggtgaggaactaaac

20 **SEQ ID NO: 17** pAOX2ADE1glslII  
tcgaccggctgcattaatgaatcgccaacgcgcggggagagggcggttgcgattggcgctcttccgcttctcgtcac  
tgactcgtcgcgtcggctgtcggctgcggcgagcggatcagctcactcaaaggcggtaatacgggtatccacagaatca  
ggggataacgcaggaagaacatgtgagcaaaaggccagcaaaaggccaggaaccgtaaaaaggccggttgctggc  
gtttttccataggctccgccccctgacgagcatcaaaaatcgacgtcaagtcagagggtggcgaacccgacaggact  
25 ataaagataaccaggcgtttcccctggaagctccctcgtgcgtctcctgttccgacctgcccgttaccggatacctgtccg  
cctttctccctcgggaagcgtggcgctttctcatagctcacgctgtaggtatctcagttcgggtgtaggtcgttcgtccaagct  
gggctgtgtgcacgaacccccgttcagccgaccgtcgcgcttatccgtaactatcgtcttgagccaacccggtaga  
cacgacttatcgccactggcagcagccactggtaacaggattagcagagcgaggtatgtaggcggtgtacagagttcttg  
aagtgggtggcctaactacggctacactagaaggacagtatttggatctgcgctctgctgaagccagttaccttcgaaaaag

agttggtagctcttgatccggcaaaaccaccgctggtagcgggtgtttttgttgcaagcagcagattacgcgcagaa  
 aaaaaggatctcaagaagatcctttgatctttctacggggtctgacgctcagtggaacgaaaactcacgttaagggattttgg  
 tcatgagattatcaaaaaggatcttcacctagatcctttaaattaaaaatgaagttttaatcaatctaaagtatatatgagtaaa  
 cttggtctgacagttaccaatgcttaatcagtgaggcacctatctcagcgatctgtctatttcgttcatccatagttgcctgactcc  
 5 ccgctggtgtagataactacgatacgggaggggttaccatctgccccagtgctgcaatgataccgcgagacccacgctcac  
 cggctccagatttatcagcaataaaccagccagccggaaggggcggagcgagaagtggctcctgcaactttatccgctcca  
 tccagctctattaattgttgcgggaagctagagtaagtagttgccagttaatagtttgcgcaacgttgttgccattgctacagg  
 catcgtggtgtcacgctcgtctgttggtatggcttcattcagctccggttccaacgatcaaggcgagttacatgatccccat  
 gttgtgcaaaaaagcggtagctccttcggtcctccgacgttgtcagaagtaagttggccgagtggtatcactcatggttatg  
 10 gcagcactgcataattcttactgtcatgccatccgtaagatgcttttctgtgactggtgagtactcaaccaagtcattctgaga  
 atagtgtatgcggcgaccgagttgctcttgcggcgtaatacgggataataccgcgccacatagcagaactttaaagtgc  
 ctcatcattggaaaacgttctcggggcgaaaactctcaaggatctaccgctgttgagatccagttcgtatgaacccactcgt  
 gcacccaactgatcttcagcatcttttactttaccagcgtttctgggtgagcaaaaacaggaaggcaaaatgccgcaaaaaa  
 ggggaataaggggcgacacggaaatgttgaatactcatactcttcttttcaatagctccaaggcaacaattgactactcagac  
 15 cgacattcattcgttattgattttaaataacgataaacggaatggttacttgaatgatttcattttatgatcattgttactaattacc  
 taaataggattttatatggaattggaagaataagggaatttcagatgtctgaaaaaggcgaggagggtactaatcattcaag  
 cccatttcttgcagtaattgcttcataagcttcaatatacttttcttactcttgatagcaatttctgcatccatggctacgccctctt  
 tggcattcaatccggtggcgtcaaccaatctctgagaaaactgcttatcgtactcttctgcgatttaccacttggttaagctttt  
 gattccaaaatctagaagaatctggagttaaaacttcactactagtagccaattcattgtttcgtccagtccaaattcgaatttcgt  
 20 atcagcaataatgatcccttcaaaaggcggaagtttttcgagcagaatacaactcgaccgccttgacagcgaccttctcac  
 aaatgtctttacctacaatctcagcagcttgttcaatagagatgtttcatcgtgttcaccctgttcagctttcgttgaaggtgtgaa  
 aatcgaggttggaaggcgtcgtctcttgaaggttctcgtttcaaccttgactccatggacagttttgagttcttgtactcttc  
 catgcacttccagtgtatgtaacctctgacaatggcttccaaagggtatcagctctgtgctttttactatcaaggatcgtccctcta  
 tgagatttgtatttttctcagacagttttgatggtagtaaagcaaaacttctgtcattagaagcaacaaatgattctttatgta  
 25 ggggtccaaaaaatcaaacagaaaactgagagctgagtgcaaaatcttcccttatcaggaataaccgttgtcataatcacatc  
 gtaagcgggagatacggtcagttgcgacgaacagcaagttgttctcatcgactgcataaatgtcttaacctttccttggcgatt  
 aaaggtaggattccgtccagatcagttgcacaatggacatacttgaaggatacagcaaaagtgtgttgaagcgatgacac  
 atggaaggaatttttcagtttctagtagtatattggggcgggtgaaagttcagatgtttaatgcttaatactcttatactcttc  
 aaagcggccaagtgttctgccaacctgactttttctgaataatgaatcgttcaagtggagtatttaaaccatgattaagttacgt

gatttggcactggataaggtcgaaaaatatccgtattcataaacgattattggtaaaagtacaaaataccactaattacggag  
aagcttagtaaacagtatcatctcttggcgttaaacgcttacaatttccattcgcattcaggctgcgcaactgttgggaaggg  
cgatcgggtgcgggcctcttcgtattacgccagggcctcgaggcacaaacgaacgtctcacttaatcttctgtactctgaaga  
ggagtgggaaataccaagaaaaacatcaaacctgaatgattttcccaaacccctaccacaagatattcatcagctgcgagat  
5 aggctgatcaggagcaagctcgtacgagaagaaacaaaatgacaaaaaaatcctatactatataaggttacaataaaaaa  
gtatcaaaaaatgaagcctgcatctctcaggcaaatggcattctgacatcctcttgattagaatctagcaagaccggcttctcgt  
aagtgcccaactgaactgaggaacagtcgtctaaggctaaaactcaatgatgatgatgatgatggcgcacggcgctattc  
agatcctcttctgagatgagttttgtcgggccccaaaaataacttcccaatcttcagttatgtcaagcgataggtaagaatggt  
gatcttattatcattctcatatgacgatctagtcggcaatgaactttctcgcatttttttgactttaatactgtctttcaacggtatgt  
10 tgtgttgtaaattattcttgcgataatgatttttcaatattggattcctcaaaagtattgtgtgaatgcctgtcaaattctcgggaa  
tatgacttcgaacatttttaaggtattgtttcgaaaaatgaactgagtttctacgtactcacctcttggtagccaaaagtttctcca  
tcatcaacataaagatctccaacggctcgtccctcggtatcaggggctataactattacatatggatcgttttcattaacattga  
agatcttctatacttattcttcatagtataatgtgaccgccttcaataaataatggaattttatccaatggtgcagaaatattcttt  
ctatcaaatcagtaccattgttataaaagagtgtaaagatgcgaattcatagaatataccgggtgggaaaaccatttccgttct  
15 gattgaccaggctccgtgacaggttgactaataagacctgaattactccagtaaaattggtatcgatatgatacaattcagcaa  
attcagggtgttcaataaacattggattcattatcggaatccagtgacacttgattatgaacatggtgtataaggtaggtag  
caggaaatatctcaattggataatatcactgactatcgacttcaaagggttcattgaataagtatggttctcttcttgggtgtctata  
tgggcgtgtgtcttaaaaaatgggtaccataagcccgttggtaccaacgtgcaatcaattcaggtgtaggtcctcagcaa  
agccagctatgtcggctctataaatggcataccagcaatgttggttgacagaaccataggaatggaaatctttaagtaatccc  
20 aattggccacattgtcaccagtcctatgtggcagcagtagcttgagagccggcaaaaaagcccttgttagaaggaaaggac  
gcttatcggatgtgaataaatcgattttattgcgtcgttaagtagcttcatgactgatagaccatatatgttatggacggatcttt  
cctcaatgtaattgtcgtgaatcaaatcttttgagctgtggtctctgggccatcgaaaatcgagggtcgttcatatcattccaa  
atgaataaattagttaaatcagccggcagatccataaacggttcgaaaaaggacttcaaactctttggccatatttgcttatggt  
atcaatccatatagaattacctggccagcaatgacctacatagtcatttccattgtgatccttgactgctacatttccattaattacc  
25 ctgtcactgatttcataatcttctttaaatgaggatcgattagtagcacaagatttctacccaactttttaattggataacagcct  
ttttggattgggaaaggagtgtgcttccaagtaaaatattttgtcgttcgtatactccaagtccaacaaataaaatcgtaag  
gaatcatatgagcatccatctgagagtcactgtgagaacgtccatctcatcattataattccatctacattgatgttacctata  
gaggaaatggcggtaaaaagggtctaccagtcaaatcggtaaattgtcaatgatagttggaatatctggccccaggggaca  
tgactacatctatgacaccatttctggagatccaatgagtcatcgttttattttactggtgtcatactttatgtctaccaagtgtca

gctgcattgacccaaaagatagatgtggacgaagatgaaaacatgaatgggatcgaaccgtacattggttggtggtaccg  
atgtgtactcaaagacatcaacgttgaaaagcctgtagggttcctttccacctgaagtgtccatcagccttagcgacgtcgca  
tgttccggtataccgtagacattagtagaacccatgaagagaaaatctagcgcaaccgattcaggcccccaggcatagag  
tcatgctttgaatacaagaaattgtccttaaacatgttgaaggtgttcttctggcagcacgtgtgcgaagtttctgcttagtgc  
5 tatgatgttcaatgttcaggaaatttgcctgttacaataagtttcagcgcatttggcagtaaacittcaattgaaaagggtcagc  
aaagatttctacggatacatcaccgttcgaagatgaaatgtgtctgcagtgagggttgaaagtgacaaaaatgaagatatttgc  
gaccagaatgagttcacagttgttttgcctaagggaagtgggaattgtggaatactggtcctgttcgctcctcttgaaatttcttg  
cgaatgcgtacttccaggtctcattgaaccgttgtgaagagatcaacaaccgctgctgttggtggcattctcttctctcattt  
atagtgaaccttactgagtgatcctgtaaaaagagagagagaatgggaactgaacggctatatcatcgccctccaattcttg  
10 gtatagttttaattatggtagcatgaagcacattcttaaggtcgtgtgcaatagactcggcgtccacttgaatagcagtg  
atgagatttggcaatatttctgcataaacctgttctatggcaaaaccagattgcgcacacttcttaatagatagtcggtaa  
acgcatgcgaaaaagcggtaaagaagaccaattggcatacagaccatttcaaaaggaccatctcgaggtaccgatccgag  
acggccggctgggccacgtgaattctttctcagttgattgtttgtgggatttagtaagtcgtaaacttcgttaaaaaagatca  
atgtagtcaatacagttgatccgaaatagaagggaagggttgcaatgtgtaagaacaatgtagttaaagcccgttttaagac  
15 aatattcttgatgctgatcagaaaaggacaataagggttttgggttcttctttataccaataatcgtctcctcatcgcttaattt  
ctccccatctcaaccggtgaagggtaggacgcttctgtaactgttcacataaaagggttttactccgagacaaaaatttat  
gcgacaaaaatagcctatcttgaagggtgatgtcttatcaactgcattgttgcaaggagaagcaaggacaactcaacatgg  
gtaaaaattcaaaaccaaccaattggaaactcccaactgtccactaggtagctgacagctgtcacttttgcgttcgttgcctgt  
ctctttcgcttaactgtcctttacggcttctttccaaaaaatcattggggaaatgtgccctcatcagagtccaatgacccat  
20 gaataaagtttctgtactgttaagacgatgaattgcaacgataatccgagcagtttacgggtacatcacgtgctttgcata  
gatctcggagtcggatcagttccggatgtgatgtattaccccatagtttcaactctaatagcagccgccaagtgcatacacc  
ctccatcaatctatgcttaagttttcaccatcgttgggtggtgatgatgactcgttagtctctgctgttcgatattaactttgtaa  
ggatcgccttggatggaaaattgaggggttgaacctgaattgcaggctacttacattggactttgagaaggctggacgg  
ttgatgaagagggctgggtgcagaggaatggaaaaaatttagttgagaggactgcttgaattttaggaaatggagtccttt  
25 aagctgacaaaactcaaggatggggattttcatgtagcttttcatgccttcgacaagctaaaggaaggaattgattctggat  
aaatggatatttgatctgcttagcagatgtcaaagtctactagtatgactcgttatctcgtagccttcaattgggcgtatctta  
ctcgaagtgttatatttttagctgacgagacgaagaacgagagagtattgacacattcagaggtaagacaatatgtcgtattat  
caaaaataagtatcgaacctctattaggagcctactggctcaaatgtgcaaccttagtggtgattgtctctgcttctgatcacaat  
ctgtcgtgtttgagagtgccgatgtatgatttttagtaaatgttttcagaaaaggcgctaagtaataaccagtaagtaataaat

aacgtaaaagtattgaatcataaaagaatcaagatagagggtcaaagcatagataatcccccgtagaagatggaccggtc  
atatgggtctgaaaaaagatctgatctcatg

SEQ ID NO: 18 pGAPZAGLSII

5 tcgagatggctctttgaaatggctcgtatgccaatgggtcttctttaccgcttttcgcatgcgttaccgactatctattaaagaa  
gtgtgcgcaatctgggtttgccatagaacagggttatgcagaaaattgccaaatctcatcactgctattacaaagtgga  
cgccgagtcattgcacacgatcctttagagaatgtgcttcatgctaccataattaaaactataccaagattggagggcgatga  
tatagccgttcagttcccattctctctctctttttacaggatcactcagtaagggtcactataaatgagaaagagagaatgccaa  
ccaacagcagcgggttgatctcttcacaacgggtcaatgagacctggaagtacgcattcgacaagaaattcaagagga  
10 ggcgaaacaggaccagtattccacaattccacttcctaagcaaaaacaaactgtgaactcattctggtcgaaaatatcttcattt  
ttgtcactttcaaactccactgcagacacattcatcttcgaaacgggtgatgtatccgtagaaatctttgctgaacctttcaattg  
aaagtftactggcaaatgcgctgaaacttattgtaaacgagcaaaattcctgaacattgaacatcatagaactaagcagga  
aaacttcgcacacgtgctgccagaagaacaactttcaacatgttaaggacaattcttgtattcaagcatgactctatgcctt  
tggggcctgaatcggttgcgctagatttctttcatgggttctactaatgtctacgggtataccggaacatgcgacgtcgctaag  
15 gctgatggacacttcaggtggaaggaacccacaggttttaacgttgatgtctttgagtacaacatcggtaccagccaac  
caatgtacgggtcgtatccattcatgtttcatcttcgtccacatctatctttgggtcaatgcagctgacacttgggtagacataa  
agtatgacaccagtaaaaaataaacgatgactcattggatctccgaaaatgggtcatagatgtatgtatgccttggggcca  
gatattccaactatcattgacaaattaccgatttgactggtagacccttttaccgcccatttctctatagggtaccatcaatgta  
gatggaattataatgatgagatggacgttctcacagtggactctcagatggatgctcatatgattccttacgattttattgggtgg  
20 acttggagtatacgaacgacaaaaatattttacttggagcagcactccttcccaatccaaaaaggctgttatccaaattaa  
aaagttgggtagaaatcttgtcgtactaatcgtatcctcatttaagaaagattatgaaatcagtgcagggtaattaatgaaat  
gtagcagtcaaggatcacatggaatgactatgtaggtcattgctggccaggtaattctatatggattgataccataagcaaa  
tatggccaaaagatttgaagtccttttcgaacgggttatggatctgccggctgatttaactaatttattcatttgggaatgatatga  
acgagccttcgattttgatggccagagaccacagctccaaaagatttgattcacgacaattacattgaggaaagatccgtc  
25 cataacatatatgggtctatcagtgcataagctacttacgacgcaataaaatcgatttaccatccgataagcgtccttctt  
tctaacaagggttttttgcgggtctcaacgtactgctgccacatggactgggtgacaatgtggccaattgggattacttaag  
atttccattcctatggttctgtcaacaacattgctgggtatgccatttataggagccgacatagctggcttggctgaggatcctac  
acctgaattgattgcacgttggtaccaagcgggcttatgggtaccatttttagagcacacgcccatatagacaccaagagaa  
gagaaccatacttattcaatgaaccttgaagtcgatagtcagtgtatattatccaattgagatatttctgctacctaccttataca

ccatgtttcataaatcaagtgtcactggatttccgataatgaatccaatgtttattgaacacctgaatttgcgaattgtatcatat  
 cgataaccaattttactggagtaattcaggtctattagtcacacgtgtcacggagcctggtaatcagaaacggaaatggtttc  
 ccacccggtataattctatgaattcgcactctttacactctttataaacaatggtactgattgatagaaaagaatattctgcacat  
 tggataaaattccattatttgaaggcgggtcacattatcactatgaaagataagtagaagatcttcaatgtaatgaaaaac  
 5 gatccatataataagttatagcccctgataccgagggacgagccgttggagatctttatgttgatgatggagaaacttttggct  
 accaaagagggtgagtacgtagaaactcagttcatttgcgaaacaataccttaaaaaatgttcgaagtcattatcccagaaattt  
 gacaggcattcaccacaatactttgaggaataccaatattgaaaaaatcattatcgaaagaataattacaacacaacataac  
 gttgaaagacagtattaaagtcacaaaaaatggcgaagaaagttcattgccgactagatcgtcatatgagaatgataataaga  
 tcaccattctaacctatcgcttgacataactgaagattgggaagttattttgggcccgaacaaaaactcatctcagaagagg  
 10 atctgaatagcgccgtcgaccatcatcatcatcattagtttagccttagacatgactgttcctcagttcaagttgggcact  
 tacgagaagaccggtcttctagattctaataagaggatgacagaatgccatttgcctgagagatgcaggcttcattttgata  
 ctttttatttgaacctatatagtataggttttttgcattttgttcttctcgtacgagcttgcctgatcagcctatctcgcagct  
 gatgaatatcttggtaggggttgggaaaatcattcgagtttgatgttttcttggattttccactcctctcagagtacagaa  
 gattaagttagaccttctgttgcggatccccacacaccatagcttcaaaatgttctactcctttttactcttcagattttctc  
 15 ggactccgcgcacgcctgaccacttcaaacacccaagcacagcactaaattttccctcttcttctctaggggtgcgtta  
 attaccgactaaagggttggaaaagaaaaagagaccgcctcgtttcttttctcgtcgaaaaaggcaataaaaattttatc  
 acgttcttttcttgaaatttttttttagtttttcttctcagtgacctccattgatatttaagttataaacggcttcaatttctcaa  
 gtttcagtttcttttctgttctattacaacttttttacttctgttcattagaaagaaagcatagcaatctaataaggcggtgt  
 tgacaattaatcatcggcatagtatatacggcatagtataatacgacaagggtgaggaactaaaccatggccaagttgaccagt  
 20 ccgttcgggtgctcaccgcgcgcgacgtcgccggagcgggtcgagttctggaccgaccggctcgggttctccgggacttc  
 gtggaggacgacttcgcccgtgtgtcgggacgacgtgacctgttcacagcgcggtccaggaccaggtgtgtccgg  
 acaacaccctggcctgggtgtgggtgcgcggcctggacgagctgtacgccgagtggtcggaggtcgtgtccacgaacttc  
 cgggacgcctccgggcccggcatgaccgagatcggcgagcagccgtgggggagggttcgcctgcgcgacccgg  
 ccggcaactgcgtgcacttcgtggccgaggagcaggactgacacgtccgacggcgccacgggtccaggcctcgga  
 25 gatccgtcccccttttcttctgtcgatatcatgtaattagttatgtcacgcttacattcacgccctccccccacatccgctctaacc  
 gaaaaggaaggagttagacaacctgaagtctaggtccctatttttttatagttatgtagtattaagaacgttatttatattca  
 aattttcttttttctgtacagacgcgtgtacgcatgtaacattatactgaaaaccttgcctgagaagggtttgggacgctcgaa  
 ggctttaatttgcaagctggagaccaacatgtgagcaaaaggccagcaaaaggccaggaaccgtaaaaaggccgcgttg  
 ctggcgttttccatagggtccgccccctgacgagcatcacaaaaatcgacgctcaagtcagagggtggcgaaacccgaca



ggactataaagataaccaggcggtttccccctggaagctccctcgtgcgctctcctgttccgaccctgccgcttacggatacct  
 gtccgcctttctcccttcgggaagcgtggcgctttctcaatgctcacgctgtaggtatctcagttcgggtgtaggtcgttcgctcc  
 aagctgggctgtgtgcacgaacccccgttcagcccaccgctgcgccttatccggtaactatcgtcttgagtccaacccgg  
 taagacacgacttatcgccactggcagcagccactggaacaggattagcagagcgaggtatgtaggcgggtgctacagag  
 5 ttctgaagtgggtggcctaactacggctacactagaaggacagtatttggtatctgcgctctgctgaagccagttaccttcgga  
 aaaagagttgtagctcttgatccggcaacaaaccaccgctggtagcgggtgtttttgttgcaagcagcagattacgcg  
 cagaaaaaaggatctcaagaagatcctttgatctttctacggggctgacgctcagtggaacgaaaactcacgttaaggg  
 attttggtcatgcatgagatcagatctttttgtagaatgtcttggtgtcctcgtccaatcaggtagccatctctgaaatatctgg  
 ctccgttgcaactccgaacgacctgctggcaacgtaaaattctccgggtaaaacttaaatgtggagtaatggaaccagaaa  
 10 cgtctcttcccttctctccttccaccgccggtaccgtccctaggaattttactctgctggagagcttcttctacggccccctt  
 gcagcaatgctcttccagcattacgttgcgggtaaaacggaggtcgtgtacccgacctagcagcccagggtggaagaaag  
 tcccggccgtcgtggcaataatagcgggcggacgcatgtcatgagattattggaaccaccagaatcgaatataaaaggc  
 gaacacctttccaattttggttctcctgacccaaagactttaaatattattttgtccctatttcaatcaatgaacaactatttc  
 gaaacgagggaattcacgtggcccagccggcgtctcggatcggtacc

15

**SEQ ID NO: 19 pGAPADE1 glsII**

tcgaccggctgcattaatgaatcgccaacgcgcggggagaggcgggttgcgtattgggcgctcttccgcttctcgtcac  
 tgactcgtcgcgtcggctgttcggctgcggcgagcgggtatcagctcactcaaaggcggtaatacgggtatccacagaatca  
 ggggataacgcaggaagaacatgtgagcaaaaggccagcaaaaggccaggaaccgtaaaaaggccgcgttgctggc  
 20 gttttccataggctccgccccctgacgagcatcacaaaaatcgacgctcaagtcagagtggtggcgaacccgacaggact  
 ataaagataaccaggcggtttccccctggaagctccctcgtgcgctctcctgttccgaccctgccgcttaccggatacctgtccg  
 cctttctcccttcgggaagcgtggcgctttctcatagctcacgctgtaggtatctcagttcgggtgtaggtcgttcgtccaagct  
 gggctgtgtgcacgaacccccgttcagcccaccgctgcgccttatccggtaactatcgtcttgagtccaacccggtgaaga  
 cacgacttatcgccactggcagcagccactggaacaggattagcagagcgaggtatgtaggcgggtgctacagagttcttg  
 25 aagtgggtggcctaactacggctacactagaaggacagtatttggtatctgcgctctgctgaagccagttaccttcgaaaaag  
 agttggtagctcttgatccggcaacaaaccaccgctggtagcgggtgtttttgttgcaagcagcagattacgcgcagaa  
 aaaaaggatctcaagaagatcctttgatctttctacggggctgacgctcagtggaacgaaaactcacgttaagggattttgg  
 tcatgagattatcaaaaaggatcttcacctagatccttttaaaataaaatgaagtttaaatcaatctaaagtatatatagtaaa  
 ctgggtctgacagttaccaatgcttaatcagtgaggcacctatctcagcgatctgtctatttcgttcatccatagttgcctgactcc

ccgtcgtgtagataactacgatacgggagggcttaccatctggccccagtgtgcaatgataccgcgagacccacgtcac  
cggctccagatttatcagcaataaaccagccagccggaaggggccgagcgcagaagtggctctgcaactttatccgcctcca  
tccagtctattaattgttgccgggaagctagagtaagtagttgccagttaatagtttgccaacgttggtgacattgctacagg  
catcgtggtgtcacgctcgtctgttggtatggcttcattcagctccgggtcccaacgatcaaggcgagttacatgatccccat  
5 gttgtgcaaaaaagcggtagctccttcggctcctccgatcgtgtgcagaagtaagtggccgcagtggtatcactcatggttatg  
gcagcactgcataattcttactgtcatgccatccgtaagatcgtttctgtgactggtgagtactcaaccaagtcattctgaga  
atagtgtatgcggcgaccgagttgctcttggccggcgtcaatacgggataataccgcgccacatagcagaactttaaaagt  
ctcatcattggaaaacgttctcggggcgaaaactctcaaggatcttaccgctgttgagatccagttcgatgtaacccactcgt  
gcacccaactgatcttcagcatctttactttcaccagcgtttctgggtgagcaaaaacaggaaggcaaaatgccgcaaaaa  
10 gggataaaggcgacacggaaatgtgaatactcatactcttcttttaataagctccaaggcaacaattgactactcagac  
cgacattcattcgttattgatitaaatcaacgataaacggaatggttactgaatgatttactttatgatcattgtttactaattacc  
taaataggattttataggaaatggaagaataagggaatttcagatgtctgaaaaaggcgaggagggtactaatcattcaag  
ccatttcttgccagtaattgcttcataagcttcaatacttttcttactcttgatagcaatttctgcacccatggctacgccctctt  
tgccattcaatccgttgccgtcaaccaatctctgagaaactgcttctgtaactctcttgcgatttaccacttggttaagtcttt  
15 gattccaaaatctagaagaatctggagttaaaactcactactagtagccaattcattgtttcgtccagtcctaaatcgaatttcgt  
atcagcaataatgatcccttcaaaaggcggaagtttttcgagcagaatacaactcgaccgccttgacagcgaccttctcac  
aaatgtctttacctacaatctcagcagcttggtcaatagagatgttttcacgtgttcaccctgttcagctttcgtgaagggtgaa  
aatcgaggttgaaaggcgtcgtctcttgaagggttcgtttcaaccttgactccatggacagttttgagttcttgactcttctc  
catgcacttccagtgtatgaacctctgacaatggcttccaaagggtatcagctctgtgctttttactatcaaggatcgtccctctaat  
20 tgagatttgatttttctcagacagtttgatggtagtaaagcaaaagacttcttgcattagaagcaacaaatgattctttatgta  
gggtgccaaaaatcaaccagaaaactgagagctgagtgcaaaatctttcccttatcaggaataaccgtttgtcataatcacatc  
gtaagcggagatacggtagtgcgacgaacagcaagttgttctcatcgactgcataaatgtctctaacctttcctttggcgatt  
aaaggtaggattccgtccagatcagtggtcacatggacatacttgaaggatacagcaaaagtgtgtggaagcgatgacac  
atggaaagggaattttcaggttcttagagtagtatattggggcggtgaaagttcagatgttaatgcttaatactcttatactcttc  
25 aaagcgcctcaagtgttctgccaacctgactttttctgaataatgaatcgttcaagtggagttttaaacatgattaagttacgt  
gatttggcactggataaggtcgaaaaatatccgtattcataaacgattattggtaaaagttacaaaataaccactaattacggag  
aagcttagtaacagttatcatctcttggtcgattaacgttacaatttccattcgcattcagggtgcgcaactgttggaagg  
cgatcgggtgcgggcctctcgtattacgccagggcctcgaggcacaacgaacgtctacttaattctctgtactctgaaga  
ggagtgggaataccaagaaaaacatcaaaactgaatgattttcccaaacccctaccacaagatattcatcagctgcgagat

aggctgatcaggagcaagctcgtacgagaagaacaaaatgacaaaaaaatcctatactatataggttacaaataaaaa  
gtatcaaaaatgaagcctgcatctctcaggcaaatggcattctgacatcctcttgattagaatctagcaagaccggctctctgt  
aagtgcccaacttgaaactgaggaacagtcagcttaaggctaaaactcaatgatgatgatgatggcgacggcgctattc  
agatcctctctgagatgagttttgtcgggccccaaaaataacttcccaatcttcagttatgtcaagcgatagggttaagaatggt  
5 gatcttattatcattctcatatgacgatctagtcggcaatgaactttctcgccattttttgactttaatactgtcttcaacggtatgt  
tgtgttgtaaattattcttgcgataatgatttttcaatattggtattcctcaaggtattgtggtgaatgcctgtcaaatctcgggaa  
tatgacttcgaacatttttaagggtattgtttcgaaaaatgaactgagtttctacgtactcacctcttggtagccaaaagtctcca  
tcatcaacataaagatctccaacggctcgtccctcggtatcaggggctataactattacatatggatcgttttcattaacattga  
agatctctatacttattcttcatagtataatgtgaccgcttcaataataatggaattttatccaatgggtgcagaaatattcttt  
10 ctatcaaatcagtagcattgttataaaagagtgtaaagatgcgaattcatagaatataccgggtgggaaaaccatttccgtttc  
gattgaccagggtccgtgacagggttgactaataagacctgaattactccagtaaaattggtatcgatatgataaattcagcaa  
attcagggtgttcaataaacattggattcattatcggaatccagtgacacttgattatgaacatgggtgtataaggtaggtag  
caggaaatatctcaattggataatatcactatcgacttcaaagggttcattgaataagtatggtctcttcttgggtgtctata  
tgggcgtgtgctctaaaaatgggtaccataagcccgttggtaccaacgtgcaatcaattcagggtgtaggatcctcagcaa  
15 agccagctatgtcggctctataaatggcataccagcaatgttgttgacagaaccataggaatggaaatctttaagtaatccc  
aattggccacattgtcaccagtcctatgtggcagcagtagcttgagagccggcaaaaaagcccttggtagaaggaaaggac  
gcttatcggatggtgaataaatcgattttattgcgtcgttaagtagcttcatgcactgatagaccatatatgttatggacggatcttt  
cctcaatgtaattgtcgtgaatcaaatcttttgagctgtggtctctgggccatcgaaaatcgagggtcgttcatatcattccaa  
atgaataaattagttaaatcagccggcagatccataaacggttcgaaaaaggacttccaaatctttggccatatttgcttatggt  
20 atcaatccatatagaattacctggccagcaatgacctacatagtcatttccattgtgatccttgactgctacattttcattaattacc  
ctgtcactgatttcataatctttcttaaatgaggatcgattagtagcacaagatttctaccaactttttaattggataacagcct  
ttttgattgggaaaggagtgctgcttccaagtaaaatattttgtcgttcgtatactccaagtccaacaaataaaatcgtaag  
gaatcatatgagcatccatctgagagtcactgtgagaacgtccatctcatcattataattccatctacattgatgttacctata  
gaggaaatggcggttaaaaagggtctaccagtc aaatcggtaaattgtcaatgatagttggaatatctggccccagggaca  
25 tgactacatctatgacaccatttctggagatccaatgagtcacgttttatttttactgggtgcatactttatgtctaccaagtgtca  
gctgcattgacccaaaagatagatgtggacgaagatgaaaacatgaatgggatcgaaccgtacattggttggtggtaccg  
atgttgactcaaagacatcaacgttgaaaagcctgtagggttcttccacctgaagtgtccatcagccttagcgacgtcgca  
tgttccggtataccgtagacattagtagaacccatgaaagagaaatctagcgcaaccgattcaggcccaaggcatagag  
tcatgctttgaatacaagaaattgccttaaacatgttgaaagttgttcttctggcagcacgtgtgcgaagtttctgcttagttc

tatgatgtcaatgttcaggaaattttgctcgtttacaataagtttcagcgcattttgccagtaaactttcaattgaaaagggtcagc  
 aaagatttctacggatacatcaccgtttcgaagatgaaatgtgtctgcagtgaggatttgaaagtacaaaaatgaagatatttc  
 gaccagaatgagttcacagttgttttgcctaaggaaagtggaattgtgaatactggcctgttcgcctcctcttgaaatttctgt  
 cgaatgcgtacttccaggtctcattgaaccgttgtgaagagatcaacaaaccgctgctgttggttgacattctctttctcattt  
 5 atagtgaaccttactgagtgatcctgtaaaaagagagagagaatgggaactgaacggctatatcatcgccctccaatcttg  
 gtatagtttaattatggtagcatgaagcacattctctaaaggatcgtgtgcaatagactcggcgtccactttgtaatagcagtg  
 atgagatttggaatattttctgcataaacctgtttctatggcaaaaccagattgcgcacacttcttaatagatagtcggtaa  
 acgcatgcgaaaaagcggtaagaagaccaattggcatacagaccatttcaaaaggaccatctcgaggtaccgatccgag  
 acggccggctgggccacgtgaattcctcgtttcgaaatagttgttcaattgattgaaatagggacaaaataaattaaattaaagt  
 10 ctttgggtcaggagaaacaaaattgggaaaggtgttcgcctttatattcgattctggtggtttccaataatctcatgacatgcg  
 tccgcccgtattattgccagcgacggccgggactttccatccctgggctgctaggtcgggtacacgacctccgtttacc  
 gcaacgtaatgctgggaagagcattgctgcaagggggccgtagaagaagctctccagcagagtaaaatttctaggggacg  
 gtaacgggcgggtggaaggagagagaagggaagagacgtttctggttcattactccacatttaagtttaccgggagaattt  
 tacgttgcagcaggtcgttcggagttgcaacggagccagatatttcagagatggctacctgattggacgaggacaccaag  
 15 acatttctacaaaaagatctgatctca

**SEQ ID NO: 20 pPICZAGLSII**

cgaacaaaaactcatctcagaagaggatctgaatagcggcgtcgaccatcatcatcatcatcattgagttgtagccttagac  
 atgactgttcctcagttcaagttgggcacttacgagaagaccggctcttgctagattctaataagaggatgtcagaatgccattt  
 20 gcctgagagatgcaggcttcattttgatactttttatttgaacctatatagtataggattttttgtcattttgttctctctgtacga  
 gcttgctcctgatcagcctatctcgcagctgatgaatatcttggtgtaggggtttgggaaaatcattcgagtttgatgttttcttg  
 gtatttccactcctcttcagagtacagaagattaagtgagaccttctgttggtcgatccccacacaccatagcttcaaaatg  
 ttttactcctttttactcttcagattttctcgactccgcgcacgcgtaccacttcaaaacaccaagcacagcatactaa  
 atttccctcttcttctctaggggtgcgttaattaccgactactaaagggttgaaaagaaaaagagaccgctcgtttcttttc  
 25 ttctgcgaaaaaggcaataaaaattttatcacgtttcttttcttgaaatttttttttagtttttcttcttcagtgacctccattgatat  
 ttaagtaataaacgggtcttcaatttctcaagtttcagtttcattttctgttctattacaacttttttacttctgttcattagaaagaaa  
 gcatagcaatctaataaggggagggtgttgacaattaatcatcgcatagtatatcgcatagtagtataacgacaaggtgag  
 gaactaaacatggccaagtgaccagtgccgttccgggtgtcaccgcgcgcgacgtcgccggagcggtcgagttctgga  
 ccgaccggctcgggttctccgggacttctgtggagacgacttcccggtgtgtccgggacgacgtgacctgttcatca

gcgcgggtccaggaccaggtggtgccggacaacaccctggcctgggtgtgggtgcgcggcctggacgagctgtacgccg  
agtggctcgaggtcgtgtccacgaactccgggacgcctccgggcccggccatgaccgagatcgcgagcagccgtggg  
ggcgggagttcgccctgcgcgacccggccggcaactgcgtgcacttcgtggccgaggagcaggactgacacgtccgac  
ggcggccacgggtcccaggcctcgagatccgtcccccttttcttctgtcgatcatgtaattagttatgtcacgcttacatt  
5 cagccctccccccacatccgctctaaccgaaaaggaaggagtagacaacctgaagtctaggtccctatttttttatagt  
tatgttagtattaagaacgttattatattcaaattttcttttttctgtacagacgcgtgtacgcatgtaacattatactgaaaacc  
ttgcttgagaaggttttgggacgctcgaaggcttaatttgaagctggagaccaacatgtgagcaaaaggccagcaaaaag  
gccaggaaccgtaaaaaggccgcgtgtgctggcgttttccataggctccgccccctgacgagcatcacaaaaatcgacgc  
tcaagtcagaggtggcgaaacccgacaggactataaagataccaggcgtttccccctggaagctccctcgtgcgctctcct  
10 gttccgacctgccgcttacggatacctgtccgcctttctcccttcgggaagcgtggcgcttttctaatgtcacgctgtagg  
tatctcagttcgggtgtaggtcgttcgctccaagctgggctgtgtgcacgaacccccgttcagcccagccgtcgccttacc  
cggtaactatcgtcttgagtcacacccggttaagacacgacttatcgccactggcagcagccactggttaacaggattagcag  
agcgaggtatgtaggcgggtgtacagagttcttgaagtgggtggcctaactacggctacactagaaggacagtatttggtatct  
gcgctctgctgaagccagttaccttcggaaaaaggttggttagctcttgatccggcaaaacaaccaccgctggtagcgggtg  
15 gttttttgttgcaagcagcagattacgcgcagaaaaaaggatctcaagaagatcctttgatcttttctacggggtctgacgc  
tcagtggaaacgaaaactcacgttaagggttttggctcatgagatcagatctaataccaagacgaaagggttgatgaaacct  
ttttccatccgacatccacaggtccattctcacacataagtgccaaacgcaacaggaggggatacactagcagcagaccg  
ttgcaaacgcaggacctccactcctcttctcctaacacccacttttgccatcgaaaaaccagcccagttattgggcttgattg  
gagctcgtcattccaattccttctattaggctactaacaccatgactttattagcctgtctatcctggccccctggcgagggttc  
20 atgtttgtttatttccgaatgcaacaagctccgcattacacccgaacatcactccagatgagggctttctgagtggtgggtcaa  
atagtttcatgttccccaaatggcccaaaactgacagtttaaacgctgtcttggaacctaatatgacaaaagcgtgatctcatcc  
aagatgaactaagtttggttcgttgaaatgctaaccggcagttgggtcaaaaagaacttcaaaagtcggcataccgtttgtct  
tgtttggtattgattgacgaatgctcaaaaataatctcattaatgcttagcgagctctctatcgttctgaaccccggtgcacct  
gtgccgaaacgcaaatggggaaacacccgcttttggatgattatgcattgtctccacattgtatgctccaagattctggtgg  
25 gaatactgctgatagcctaacgttcattgatcaaaatttaactgttctaaccctacttgacagcaatatataaacagaaggaag  
ctgccctgtcttaaacctttttttatcatcattattagcttactttcataattgcgactggttccaattgacaagcttttgatttaacg  
actttaacgacaacttgagaagatcaaaaaacaactaattatcgaaacgagggaattcacgtggcccagccggcgtctcg  
gatcggtacctcgagatggtccttttgaaatggctcgtatgccaaattggtcttcttaccgcttttgcgatgcgtttaccgactat  
ctattaaagaagtgtgcgcaatctgggttttccatagaaacagggttatgcagaaaatttgccaaatctcatcactgctatt

acaaagtggacgccgagtcattgcacacgatcccttagagaatgtgcttcacgtaccataaataaaactataccaagattgg  
agggcgatgatatagccgttcagttccattctctctctctttttacaggatcactcagtaagggttcactataaatgagaaagag  
agaatgccaaccaacagcagcgggttgatctcttcacaacgggtcaatgagacctggaagtacgcaatcgacaagaaatt  
tcaagaggaggcgaacaggaccagtattccacaattccattcccttaagcaaaaacaaactgtgaactcattctggtcgaaa  
5 atatcttcattttgtcactttcaaactccactgcagacacatttcattcttgaaacgggtgatgtatccgtagaaatctttgctgaac  
ctttcaattgaaagttactggcaaaatgcgctgaaactattgtaaacgagcaaaatttctgaacatgaacatcatagaact  
aagcaggaaaaacttcgcacacgtgctgccagaagaaacaactttcaacatgtttaaggacaatttctgtattcaaagcatga  
ctctatgccctttggggcctgaatcggttcgctagatttctcttcacgttgctactaatgtctacgggtataccggaacatgcga  
cgtcgctaaggctgatggacacttcaggtggaaaggaaccctacaggcctttcaacgttgatgtctttgagtacaacatcggt  
10 accagccaaccaatgtacgggtgatccattcatgttttcattcttcgccacatctatcttttgggtcaatgcagctgacacttg  
ggtagacataaagtatgacaccagtaaaaaataaacgatgactcattggatctccgaaaatgggtgcatagatgtagtcatgt  
ccctggggccagatattccaactatcattgacaaatttaccgattgactggtagacccttttaccgccatttctctataggg  
taccatcaatgtagatggaattataatgatgagatggacgttctcacagtggactctcagatggatgctcatatgattccttacg  
attttatttgggttgactggagtatacgaacgacaaaaaatatttacttggaaagcagcactcctttcccaatccaaaaggctg  
15 ttatccaaattaaaaaagtgggtagaaatcttgcgtactaatcgatcctcatttaagaaagattatgaaatcagtgcagggt  
aattaatgaaaatgtagcagtcaggatcacaatggaaatgactatgtaggtcattgctggccaggaattctatatggattgat  
accataagcaaatatggccaaaagatttgaagtccttttcgaacgggttatggatctgccggctgatttaactaatttattcatt  
tggaatgatatgaacgagccttcgatttgcgatggcccagagaccacagctccaaaagatttgattcacgacaattacattga  
ggaaagatccgtccataacatatatggtctatcagtgcatgaagctacttacgacgcaataaaatcgatttattcaccatccgat  
20 aagcgtcctttccttctaacaagggttttttgcggctctcaacgtactgctgccacatggactggtgacaatgtggccaatt  
gggattacttaagatttccattcctatggttctgtcaacaacattgctggatgccatttataggagccgacatagctggcttt  
gctgaggatcctacacctgaattgattgcacgttggtaccaagcgggcttatggtacccatttttagagcacacgccatata  
gacaccaagagaagagaaccatacttattcaatgaacctttgaagtcgatagtacgtgatattatccaattgagatattcctgc  
tacctaccttatacaccatgtttcataaatcaagtgtcactggatttccgataatgaatccaatgtttattgaacacctgaatttgc  
25 tgaattgtatcatatcgataaccaattttactggagtaattcaggtctattagtcaaacctgtcacggagcctggtcaatcagaaa  
cggaaatggtttccacccgggtatattctatgaattcgcacttttacactctttataaacaatggtactgatttgatagaaaaga  
atatttctgcaccattggataaaattccattatttattgaaggcggtcacattatcactatgaaagataagtatagaagatctcaa  
tgtaatatgaaaacgatccatatgtaatgttatagccctgataccgaggagcagccgttgagatctttatgttgatgatg  
gagaaacttttggctaccaaagaggtgagtacgtagaactcagttcattttcgaaaacaataaccttaaaaaatgttcgaagtc

atattcccgagaatttgacaggcattcaccacaatactttgaggaataccaatattgaaaaaatcattatcgcaagaataattt  
acaacacacataacgttgaaagacagtattaaagtcaaaaaaatggcgaagaaagttcattgccgactagatcgcatat  
gagaatgataataagatcaccattcttaacctatcgcttgacataactgaagattgggaagttattttgggcc

5 **SEQ ID NO: 21** pPICADE1glsII

aaattcctcgtttcgaataattagttgtttttgatcttctcaagttgtcgttaaaagtcgttaaaatcaaaagcttgcaattggaac  
cagtcgcaattatgaaagtaagctaataatgatgataaaaaaaagggttaagacagggcagcttccttctgtttatatattgct  
gtcaagtagggggttagaacagttaaatttgatcatgaacgttaggctatcagcagattcccaccagaatcttggaagcatac  
aatgtggagacaatgcataatcatccaaaaagcgggtgtttcccatgttcggcacaggtgcaccgggggttcagaag  
10 cgatagagagactgcgctaagcattaatgagattattttgagcattcgtcaatcaatacacaagacaaaacggatgccg  
acttttggagtttcttttgaccaactggccgtagcatttcaacgaaccaaacttagttcatcttgatgagatcacgctttgtc  
atattaggtccaagacagcgtttaaactgtcagtttgggccatttggggaacatgaaactatttgacccacactcagaaag  
ccctcatctggagtgttcgggtgtaatgcggagcttgttcattcggaaataaacaacatgaacctgccagggggggc  
caggatagacaggctaataaagtcaggtgttagtagcctaataagaaggaattggaatgagcagctccaatcaagccaat  
15 aactgggctggttttcgatggcaaaagtgggtgttgaggagaagaggagtgaggctcctgcgtttgcaacggctcgtcgtc  
agtgtatcccctcctgttgcgtttggcacttatgtgtgagaatggacctgtggatgtcggatggcaaaaagggttcattcaacct  
ttcgtctttggatgttgcgaccggctgcattaatgaatcgccaacgcgcggggagaggcgggttgcgtattgggcgctctt  
ccgcttcctcgtcactgactcgtcgcctcggtcgttcggctcggcgagcggatcagctcactcaaaggcggtaatac  
gttatccacagaatcaggggataacgcaggaaagaacatgtgagcaaaaggccagcaaaaggccaggaaccgtaaaaa  
20 ggccgcgttgctggcggttttccataggctccgccccctgacgagcatcacaaaaatcgacgctcaagtcagaggtggcg  
aaaccggacaggactataaagataccaggcgttccccctggaagctccctcgtgcgctcctgttccgacctgccgctt  
accggataacctgtccgccttctcccttcgggaagcgtggcgctttctcatagctcacgctgtaggtatctcagttcgggttag  
gtcgttcgctccaagctgggctgtgtgcacgaacccccgttcagcccagccgctgcgccttatccggtaactatcgtctga  
gtccaacccggtaagacacgacttatcgccactggcagcagccactggtaacaggattagcagagcgaggtatgtaggcg  
25 gtgctacagagttctgaagtgggtggcctaactacggctacactagaaggacagtatttggtatctgcgctcgtcgtgaagcca  
gttaccttcgaaaaagagttggtagctcttgatccggcaacaaaccaccgctggtagcgggtgtttttgtttgcaagcag  
cagattacgcgcagaaaaaaggatctcaagaagatcctttgatctttctacggggtctgacgctcagtggaaacgaaaactc  
acgtaagggaftttggtcatgagattatcaaaaaggatcttcacctagatccttttaattaaaaatgaagtttaaatcaatcta  
aagtatatatagtaaacttggctcgtacagttaccaatgcttaatcagttaggcacctatctcagcgatctgtctatttcgttcac

catagttgccctgactccccgtcgtgtagataactacgatacgggagggttaccatctggccccagtgtgcaatgataccg  
 cgagaccacgctcaccggctccagattatcagcaataaaccagccagccggaaggccgagcgcaagaagtggcctg  
 caactttatccgctccatccagtctattaattgtgccgggaagctagagtaagtagttgccagttaatagtttgcgaacgtt  
 gttgccattgctacaggcatcgtggtgtcacgctcgtcgtttggtatggcttcattcagctccggttcccaacgatcaaggcga  
 5 gttacatgatccccatgttggtgcaaaaaagcggtagctccttcggctcctccgatcgtgtcagaagtaagtggccgcagtg  
 ttatcactcatggttatggcagcactgcataattcttactgtcatgccatccgtaagatgcttttctgtgactggtgagtactca  
 accaagtcattctgagaatagtgtatcgggcgaccgagttgctcttggccggcgtcaatacgggataataccgcgccacata  
 gcagaactttaaaagtgtcatcattggaaaacgttcttcggggcgaaaactctcaaggatcttaccgctgttgagatccagtt  
 cgatgtaaccctcgtgcaccaactgatcttcagcatctttactttcaccagcgtttctgggtgagcaaaaacaggaaggc  
 10 aaaatgccgcaaaaagggaataaggcgacacggaaatgttgaatactcatactcttcttttcaatagctccaaggcaac  
 aaatgactactcagaccgacattcattcgttattgattttaaataacgataaacggaatggttacttgaatgatttcatttatga  
 tcattgtttactaattacctaataaggattttatatggaattggaagaataagggaatttcagatgtctgaaaaaggcgaggag  
 ggtactaatcattcaagcccatttcttgcagtaattgcttcataagcttcaataacttttcttactcttgatagcaatttctgcac  
 catggctacgccctcttggcattcaatccgttgccgctcaaccaatctctgagaaactgcttctgtaactctcttgcgatttac  
 15 ccacttggaagtctttgattccaaaatctagaagaatctggaggttaaaactcatctactagtaccaattcattgtttcgtccag  
 tccaaattcgaatttcgtatcagcaataatgatcccttcaaaaaggcggaagttttgcagcagaatacaactcgaccgccttg  
 acagcgacctctcacaatgtctttacctacaatctcagcagcttgttcaatagagatgtttcatcgtgttcacctgttcagct  
 ttctgtgaagggtgtgaaaatcgagttggaaaggcgctcgctctctgaagggtctcgttttcaaccttgactccatggacagttt  
 tgagttcttgactcttccatgcactccagtgtgtaacctctgacaatggcttccaaaggatcagctgtgctttttactatca  
 20 aggatcgtccctctaattgagatttgatttttcttcagacagtttgatggtagtaagcaaaagacttccttgctcattagaagcaa  
 ccaaatgattctttatgtagggtgccaaaaaatcaaacagaaaactgagagctgagtcaaaatcttcccttatcaggaatac  
 cgtttgcataatcacatcgtaaaggagatacggtcagttgcgacgaacagcaagttgttctcatcgactgcataaatgtctc  
 taaccttctttggcgattaaaggtaggattccgtccagatcagtttcacaatggacatacttgaaggatacagcaaagtg  
 tgttggaagcgatgacacatggaaaggaattttcagatttctagagtagtatattggggcggtgaaagttcagatgtttaatg  
 25 cttaatactcttatactcttcaagcgcccaagtgttctgccaacctgactttttctgaataatgaatcgttcaagtggagtattta  
 aacctgattaagttacgtgatttggcactggataaggctgaaaaatccgtattcataaacgattattggtaaaagtataaaa  
 ataccactaattacggagaagcttagtaacagttatcatcttggctgattaacgcttacaattccattcgcattcaggctgc  
 gcaactgttgggaaggcgatcgggtcggggcctctcgtattacgccagggcctcgaggcacaaacgaacgtctcactta  
 atcttctgactctgaagaggagtgggaaataccaagaaaaacatcaaaactcgaatgattttccaaaccctaccacaagat



attcatcagctgcgagataggctgatcaggagcaagctcgtacgagaagaacaaaatgacaaaaaaatcctatactatat  
agggtacaaataaaaaagtatcaaaaatgaagcctgcacatctcaggcaaatggcattctgacatcctcttgattagaatctag  
caagaccgggtctctcgttaagtcccaactgaactgaggaacagtcattgtaaggctacaaactcaatgatgatgatgatg  
atggctgacggcgctattcagatcctctctgagatgagttttgtcgggccccaaaaataactcccaatcttcagttatgtcaa  
5 gcgataggftaagaatggtgatcttattatcattctcatatgacgatctagtcggcaatgaactttctcgccattttttgacttta  
atactgtctttcaacgttatgttgtgtgtaaatattcttgcgataatgatttttcaatattggtattcctcaaagtattgtggtgaat  
gcctgtcaaattctcgggaatatgacttcgaacatttttaaggattgtttcgaanaatgaactgagttctacgtactcacctcttt  
ggtagccaaaagtttctccatcatcaacataaagatctcaacggctcgtccctcggatcaggggctataactattacatatg  
gatcgttttcattaacattgaagatcttatacttattcttcatagtgataatgtgaccgccttcaataaataatggaatttatcca  
10 atgggtgcagaaatattctttctatcaaatcagtagcattgtttataaaagagtgtaaagatgcgaattcatagaatataccgggt  
gggaaaaccatttccgttctgattgaccaggctccgtgacagggttgactaatagacctgaattactccagtaaaattggttat  
cgatatgatacaattcagcaaatcagggtgttcaataaacattggattcattatcggaatccagtgacacttgatttatgaaac  
atgggtgtataaggtaggtagcaggaaatatctcaattggataatatcacgtactatcgacttcaaagggttcattgaataagtatg  
gttctcttctcttggtgtctatatgggcgtgtgctctaaaaatgggtaccataagcccgttggtaccaacgtgcaatcaattca  
15 ggtgtaggatacctcagcaagccagctatgtcggctcctataaatggcataaccagcaatgttgttgacagaacctaggaat  
ggaaatctttaagtaatcccaattggccacattgtcaccagtcattgtggcagcagtagcttgagagccggcaaaaaaagcc  
cttgtagaaggaaaggacgcttatcggtggtgaataaatcgattttattgcgtcgttaagtagcttcatgcactgatagacat  
atatgttatggacggatcttctcaatgtaattgtcgtgaatcaaatctttggagctgtggtctctgggccatcgaaaatcgaa  
ggctcgttcataatcattccaaatgaataaattagttaaatcagccggcagatccataaaccgttcgaaaaaggacttccaaatc  
20 ttttgccatatttgcttatggtatcaatccatatagaattacctggccagcaatgacctacatagtcatttccattgtgatccttga  
ctgctacattttcattaattaccctgtcactgatttcataatcttctttaaataaggatcgatttagtacgacaagatttctaccaac  
tttttaatttgataacagcctttttgattgggaaaggagtgtgcttccaagtaaaatattttgtcgttcgtatactccaagtc  
caaccaaataaaaatcgttaaggaaatcatatgagcatccatctgagagtccactgtgagaacgtccatctcatcattataattccat  
ctacattgatgtaccctatagaggaaatgggcgttaaaaagggtctaccagtcaaatcggtaaaattgtcaatgatagttgg  
25 aatatctggccccaggacatgactacatctatgacaccatttctggagatccaatgagtcacgttttattttactggtgtcata  
cttatgtctaccaagtgtcagctgcattgacccaaaagatagatgtggacgaagatgaaaacatgaatgggatcgaaccg  
tacattggttggtgtggtaccgatgttgtactcaaagacatcaacgttgaaaagcctgtagggttcttccacctgaagtgtcca  
tcagccttagcgacgtcgcattgtccggtataccgtagacattagtagaacccatgaaagagaaatctagcgcaaccgattc  
aggcccccaggcatagagtcattgttgaatacaagaaattgtccttaaacatgttgaaagttgtttctctggcagcacgtgt

gcgaagtttctgcttagttctatgatgttcaatgttcaggaaatttgctcgttacaataagtttcagcgcattttgccagtaaa  
ctttcaattgaaaagggttcagcaaagatttctacggatacatcaccgttcgaagatgaaatgtgtctgcagtggagtttgaag  
tgacaaaaatgaagatatttgcaccagaatgagttcacagttgttttgcttaaggaagtgaattgtggaatactggtcctgt  
tcgcctcctcttgaaatttctgtcgaatgcgtactccaggtctcattgaaccgttggaagatcaacaaaccgtgctgttg  
5 gttggcattctctctttctcattatagtgaaacctactgagtgatcctgtaaaaagagagagagaatgggaactgaacggcta  
tatcatcgcctccaatcttggtatagtttaattatggtagcatgaagcacattctctaaaggatcgtgtgcaatagactcggcg  
tccacttgaatagcagtgatgagatttggcaatatttctgcataaacctgtttctatggcaaaaccagattgcgcacactt  
cttaatagatagtcggtaaacgcagtcgaaaaagcggtaaagaagaccaattggcatacagaccattcaaaaggaccatc  
tcg

10

**SEQ ID NO: 22 pYPT1ZAGLSII**

cgaacaaaaactcatctcagaagaggatctgaatagcgccgtcgaccatcatcatcatcattgagttgtagccttagac  
atgactgttcctcagttcaagttgggcacttacgagaagaccggtcttgctagattctaataagaggatgcagaatgccattt  
gcctgagagatgcaggcttcattttgatactttttatttgaacctatatagtataggattttttgtcattttgttctctcgtacga  
15 gcttgctcctgatcagcctatctcgagctgatgaatatctgtggtagggttgggaaatcattcagtttgatgttttcttg  
gtatttcccactcctctcagagtacagaagattaagtgaaccttcgtttgtcggtatccccacacaccatagcttcaaatg  
tttctactcctttttactctccagattttctcgactccgcgcacgcgtaccacttcaaacacccaagcacagcatactaa  
attttccctctttctcctctaggggtgctgtaattaccgtactaaagggttgaaaagaaaaagagaccgcctcgtttcttttc  
ttctgcgaaaaaggcaataaaaaattttatcacgtttcttttctgaaatttttttttagtttttctcttcagtacacctcattgatat  
20 ttaagttaataaacggtcttcaatttctcaagtttcagtttcatttttctgttctattacaacttttttacttctgttcattagaaagaaa  
gcatagcaatctaafctaagggcggtgttgacaattaatcatcgccatagtatatcgccatagataatacgaaggtgag  
gaactaaacctaggccaagttgaccagtgcggttccggtgctcaccgcgcgcgacgtcgccggagcggtcgagttctgga  
ccgaccggctcgggttctcccgggacttcgtggaggacgacttcgccggtgtggtccgggacgacgtgacctgttcatca  
gcgcggtccaggaccaggtggtgccggacaacacctggcctgggtgtgggtgcgcggcctggacgagctgtacgccg  
25 agtggtcggaggtcgtgtccacgaactccgggacgcctccgggcccggccatgaccgagatcgccgagcagccgtggg  
ggcgggaggtcgcctgcgcgacccggccggcaactgcgtgcacttcgtggccgaggagcaggactgacacgtccgac  
ggcggcccacgggtcccaggcctcgagatccgtccccctttcttctgcgatcatgtaattagttatgtcacgcttacatt  
cacgcctccccccacatccgtctaaccgaaaaggaaggagtagacaacctgaagtctaggtccctatttttttatagt  
tatgttagtattaagaacgttatttatatttcaatttttcttttttctgtacagacgcgtgtacgcatgtaacattatactgaaaacc

ttgcttgagaagggttttgggacgctcgaaggctttaatttgcaagctggagaccaacatgtgagcaaaaggccagcaaaag  
gccaggaaccgtaaaaaggccgcgttgctggcgttttccataggctccgccccctgacgagcatcacaaaaatcgacgc  
tcaagtcagaggtggcgaaacccgacaggactataaagataaccaggcgttccccctggaagctccctcgtgcgtctcct  
gttccgaccctgccgcttaccggatacctgtccgcctttctccctcgggaagcgtggcgttttctaatgtcacgctgtagg  
5 tatctcagttcgggtgtaggtcgttcgtccaagctgggctgtgtgcacgaacccccgttcagcccaccgctgcgccttacc  
cggtaactatcgtcttgagtcacacccggttaagacacgacttatgccactggcagcagccactggtaacaggattagcag  
agcgaggtatgtaggcgtgtacagagttctgaagtggtggcctaactacggctacactagaaggacagtatttggtatct  
gcgctctgtgaagccagttaccttcggaaaaagagttggtagctcttgatccggcaacaaccaccgctggtagcgggtg  
gtttttgttgcaagcagcagattacgcgcagaaaaaaggatctcaagaagatccttgatctttctacggggtctgacgc  
10 tcagtggaaacgaaaactcacgttaagggttttggtcatgagatcagatctatgatgagtcacaatctgcttccacagacgagt  
acaaggacaggcaaaaggaattggaagaattgctaaccaataatgagcaagttctatggagctgctggtggagctcctg  
gtggagctcctggtggcttccctggaggtttccctggcggagctggcgcagctggcgggtgccccaggtggtgctgcccc  
ggcggagacagcggaccaaccgtggaagaagtcgattaagcaattcaacggataaattctggttaatatataacgtgaat  
aggaaattaaggaaatttggatctaataatgtgctgtatgccgacatcgggcacgtagattgtatgtatcgtgacactata  
15 ataagccagccaaaaccctaaaccagttgccctccactaattagtgtactaccaatcttgctcttcgggtgtctttataag  
gacagattcacagctctgttgcccaatacacacatacacacagagataatagcagtcgaattcacgtggccagccggc  
cgtctcggatcggtagctcgtgatggtcctttgaaatggctcgtatgccaatggtcttcttaccgcttttcgcatgcgttac  
cgactatctattaaagaagtgtgcgaatctgggtttgccatagaacagggttatgcagaaaatattgccaatctcatcac  
tgctattacaaagtggacgccgagctattgcacacgatccttagagaatgtgcttcatgtaccataattaaaactataccaa  
20 gattggagggcgatgatatagccgttcagttccattctctctctttttacaggatcactcagtaagggttcactataaatgag  
aaagagagaatgccaccaacagcagcgggttggtgatctcttcacaacgggtcaatgagacctggaagtacgcattcgaca  
agaaatttcaagaggaggcgaacaggaccagtattccacaattccacttcctaagcaaaaacaaactgtgaactcattctgg  
tcgaaaatatcttcattttgtcactttcaaactccactgcagacacattcatcttcgaaacgggtgatgtatccgtagaatcttg  
ctgaacctttcaattgaaagttactggcaaaatgcgctgaaacttattgtaaacgagcaaaatttcfgaacattgaacatcat  
25 agaactaagcaggaaaacttcgcacacgtgtgccagaagaaacaactttcaacatgttaaggacaatttctgtattcaaa  
gcatgactctatgcccttggggcctgaatcggttgcgctagatttctcttcatgggttctactaatgtctacggataccggaac  
atgcgacgtcgttaaggctgatggacacttcaggtggaaaggaaacctacaggctttcaacgttgatgtctttgagtacaac  
atcggtagcagccaaccaatgtacgggtcgatcccattcatgtttcatcttcgtccacatctatctttgggtcaatgcagctga  
cacttgggtagacataaagtatgacaccagtaaaaataaacgatgactcattggatctccgaaaatggtgtcatagatgtag

tcatgtccctggggccagatatccaactatcattgacaaatttaccgatttgactggtagacccttttaccgccatttcctcta  
 tagggtagcatcaatgtagatggaattataatgatgagatggacgttctcacagtggactctcagatggatgctcatatgattcc  
 ttacgattttatttggttgacttgaggtatagcaacgacaaaaatattttacttgaagcagcactccttccaatccaaaaa  
 ggctgttatccaaattaaaaagttggtagaaatctgtcgtactaatcgatcctcattaaagaaagattatgaaatcagtac  
 5 agggtaattaatgaaaatgtagcagtcaggatcacatggaatgactatgtaggtcattgctggccaggtatctatatgg  
 attgataccataagcaaatatggccaaaagatttgaagtccttttcgaacggttatggatctgccggctgatttaactaattta  
 ttcatgtgaatgatatgaacgagccttcgatttctgatggccagagaccacagctccaaaagatttgattcacgacaattac  
 attgaggaaagatccgtccataacatatatggtctatcagtgcatgaagctacttacgacgcaataaaatcgatttattcccat  
 ccgataagcgtcctttccttctaacaagggcttttttgcggctctcaacgtactgctgccacatggactggtagaatgtggc  
 10 caattgggattacttaagatttccattcctatggttctgtcaacaacattgctggtatgccatttataggagccgacatagctg  
 gcttctgtaggacctacacctgaattgattgcacgttggtaccaagcgggcttatggtaccatttttagagcacacgccc  
 atatagacaccaagagaagagaaccatactattcaatgaaccttgaagtcgatagtacgtgatattatccaattgagatattt  
 cctgctacctacctatacaccatgttcataaatcaagtgtcactggatttccgataatgaatccaatgtttattgaacacctga  
 atttctgaattgtatcatatcgataaccaattttactggagtaattcaggtctattagcaaacctgtcacggagcctggtcaatc  
 15 agaaacggaaatggtttccaccgggtatattctatgaattcgcacttttacactctttataaacaatggtactgattgataga  
 aaagaatatttctgcaccattggataaaattccattatttattgaaggcgggtcacattatcactatgaaagataagatagaagat  
 ctcaatgttaatgaaaaacgatccatatgaatagttatagcccctgataccgaggagcagagccgttgagatctttatgttga  
 tgalggagaaacttttggctaccaaaagaggtgagtacgtagaaactcagttcattttcgaacaataccttaaaaaatgttcg  
 aagtcatttcccagagaatttgacaggcattcaccacaatactttgaggaataccaatattgaaaaatcattatcgaaagaat  
 20 aatttacaacacataacggttgaagacagtattaaagtcaaaaaaatggcgaagaaagtcattgccgactagatcgtc  
 atatgagaatgataataagatcaccattcttaacctatcgcttgacataactgaagattgggaagttatttttgggcc

**SEQ ID NO: 23** pYPT1ADE1glsII

gtcgaccggctgcattaatgaatcggccaacgcgcggggagaggcgggttgcgtattgggcgctcttccgcttcctcgctca  
 25 ctgactcgtcgcgtcggctggtcggctgcggcgagcgggtatcagctcactcaaaggcggtaatacgggtatccacagaatc  
 aggggataacgcaggaagaacatgtgagcaaaaggccagcaaaaggccaggaaccgtaaaaaggccgcttgctgg  
 cgttttccataggctccgccccctgacgagcatcacaataatcgacgctcaagtcagagggtggcgaaacccgacagga  
 ctataaagataccaggcgtttcccctggaagctccctcgtgcgctcctcgttccgacctgccgcttaccggatacctgtcc  
 gccttctcccttcgggaagcgtggcgctttctcatagctcacgctgtaggtatctcagttcgggtgtaggtcgttcgtccaagc

tgggctgtgtgcacgaacccccgttcagcccgaccgctgcgccttatccggtaactatcgctttagtccaacccggtaag  
acacgacttatcgccactggcagcagccactggtaacaggattagcagagcgaggatgtaggcgggtgtacagagtctt  
gaagtggtagcctaactacggctacactagaaggacagtatttggtatctgcgctctgctgaagccagttaccttcgaaaaa  
gagttggtagctcttgatccggcaacaacacccgctggtagcgggtggtttttgttgcaagcagcagattacgcgcaga  
5 aaaaaaggatctcaagaagatccttgatctttctacggggtctgacgctcagtggaaacgaaaactcacgttaagggttttg  
gtcatgagattatcaaaaaggatcttcacctagatcctttaaattaaaaatgaagtttaaatcaatctaaagtatatatgagtaa  
acttggctgacagttaccaatgcttaatcagtgggcacctatctcagcgatctgtctatttcgttcacatagttgcctgactc  
cccgctgctgtagataactacgatacgggaggggttaccatctggccccagtgctgcaatgataccgcgagaccacgctca  
ccggctccagatttatcagcaataaaccagccagccggaagggccgagcgcagaagtggctctgcaacttatccgcctcc  
10 atccagtccttaattgttgccgggaagctagagtaagtagttgccagttaatagtttgcgaacggttggtgccattgctacag  
gcatcgtggtgtcacgctcgtcgtttggtatggcttcattcagctccgggttcccaacgatcaaggcgagttacatgatcccca  
tgttgtcaaaaaagcggtagctccttcggctcctccgacgttggtcagaagtaagttggccgcagtggtatcactcatggttat  
ggcagcactgcataattcttactgtcatgccatccgtaagatgctttctgtgactggtgagtactcaaccaagtcattctgag  
aatagtgtatgcggcgaccgagtgctcttgccggcgctcaatacgggataataccgcgccacatagcagaactttaaaagt  
15 gctcatcattggaaaacgttcttcggggcgaaaactctcaaggatcttaccgctgttgagatccagttcagtgtaaccactcg  
tgcaccaactgatcttcagcatctttactttaccagcggttctgggtgagcaaaaacagggaaggcaaaatgccgaaaaa  
agggaafaaggcgacacggaaatgtgaatactcatactcttctttcaatagctccaaggcaacaaatgactactcaga  
ccgacattcattcggttattgattttaaatcaacgataaacggaatggttacttgaatgatttcatttatgatcattgttactaattac  
ctaaataggattttatatggaattggaagaataagggaatttcagatgtctgaaaaaggcgaggagggtactaatcattcaa  
20 gccatttcttgccagtaattgcttcataagcttcaatatactttcttactcttgatagcaattctgcatccatggctacgccctct  
ttgccattcaatccgttggccgtcaaccaatctctgagaaactgcttatcgttaactcttgcgattaccacttggttaagtcttt  
gattcaaaaatctagaagaatctggagttaaaacttcactactagtaaccaattcattgtttcgtccagtcctaaatcgaattcgt  
atcagcaataatgatcccttcaaaaggcggaagttttgcagcagaatacaactcgaccgccttgacagcgaccttctcac  
aaatgtctttacctaataatctcagcagcttgttcaatagagatgtttcactggttaccctgttcagcttctgttgaagggttgaa  
25 aatcggagttggaaggcgctcgtctcttgaagggtctctgtttcaaccttgactccatggacagttttgagttctgtactcttc  
catgcacttccagtgtatgtaacctctgacaatggcttccaaagggtatcagctgtgctttttactatcaaggatcgtccctcta  
tgagatttgatttttctcagacagtttgatggtagtaagcaagacttcttgcattagaagcaacaaatgattctttatgta  
gggtgccaaaaaatcaaacagaaaaactgagagctgagtgcaaaatctttcccttatcaggaataccgttgcataatcacatc  
gtaagcggagatacggtcagttgcgacgaacagcaagttgtctcatcgactgcataaatgtctctaacctttccttggcgatt

aaaggtaggattccgtccagatcagtggtcacaatggacatacttgaaggatacagcaaagtgtgtggaagcgatgacac  
atggaaaggaattttcaggttcctagagtagtatattggggcggtgaaagttcagatgtttaatgcttaatactcttatactcttc  
aaagcgcccaagtgttctgccaacctgactttttctgaataatgaatcgttcaagtggagtatttaaacatgattaagttacgt  
gatttggcactggataaggtcgaaaaataccgtattcataaacgattattggtaaaagttacaaaataccactaattacggag  
5 aagcttagtaacagttatcatctcttggcgtgattaacgcttacaatttcattcgccattcaggctgcgcaactgttgggaaggg  
cgatcgggtgcgggcctctcgtattacgccagggcctcgaggcacaacgaacgtctcacttaattcttctgtactctgaaga  
ggagtgggaataccaagaaaaacatcaaactcgaatgattttccaaacccctaccacaagatattcatcagctgcgagat  
aggctgatcaggagcaagctcgtacgagaagaacaaaaatgacaaaaaaaatcctatactatataaggttacaaataaaaaa  
gtatcaaaaatgaagcctgcatctctcaggcaaatggcattctgacatcctcttgattagaatctagcaagaccggtcttctcgt  
10 aagtgcccaactgaactgaggaaacagtcattgctaaaggctacaaactaatgatgatgatgatggtcgacggcgctatt  
cagatcctcttctgagatgagttttgtcgggccccaaaaataacttcccaatcttcagttatgtcaagcgatagggttaagaatgg  
tgatcttattatcattctcatatgacgatctagtcggcaatgaactttctcgcattttttgactttaatactgtctttcaacgttatg  
ttgtgtgtaaattattcttgcgataatgatttttcaatattggtattcctcaaagtattgtgtgaatgcctgtcaaattctcggga  
atatgacttcgaacatttttaaggtattgtttcgaaaatgaactgagtttctacgtactcacctcttggtagccaaaagtttctcc  
15 atcatcaacataaagatctccaacggctcgtccctcggtatcaggggctataactattacataggtatcgttttcattaacattg  
aagatcttctatacttattcttcatagtgataatgtgaccgccttcaataaataatggaattttatccaatgggtcgaaaatattcttt  
tctatcaaatcagtaccattgtttataaaagagtgtaaagatgcgaattcatagaatataccgggtgggaaaaccatttccgtttc  
tgattgaccagggtccgtgacagggttgactaatagacctgaattactccagtaaaattggtatcgcgatgatacaattcagca  
aattcagggtgttcaataaacattggattcattatcggaatccagtgacacttgattatgaacatgggtgataaggttaggta  
20 gcaggaaatatctcaattggataatcacgtactatcacttcaaagggttcattgaataagtatggttcttctcttgggtgtctat  
atgggcgtgtgctctaaaaaatgggtaccataagcccgttgggtaccaacgtgcaatcaattcagggtgtaggatcctcagca  
aagccagctatgtcggctcctataaatggcataaccagcaatgtgttgacagaaccataggaatggaaatctttaagtaatcc  
caattggccacattgtcaccagtcattgtggcagcagtagcttgagagccggcaaaaaagccctgttagaaggaaagga  
cgcttatcggatggtgaataaatcgaftttattgcgtcgtaagtagcttcatgactgatagaccatatatgttatggacggatctt  
25 tcctcaatgtaattgtcgtgaatcaaatctttggagctgtggtctctgggccatcgaaaatcgaaggctcgttcataatcattcca  
aatgaataaattagttaaatcagccggcagatccataaacggttcgaaaaaggacttccaaatctttggccatatttgcttatg  
gtatcaatccatatagaattacctggccagcaatgacctacatagtcatttccattgtgatccttgactgtacatttccattaatta  
ccctgtcactgatttcataatcttctttaaattgaggatcgatttagtacgacaagatttctaccaactttttaattggataacagc  
cttttggattgggaaaggagtgctgcttccaagtaaaatatttttgcgttcgtatactccaagtccaaccaataaaatcgtaa

ggaatcatatgagcatccatctgagagtccactgtgagaacgtccatctcatcattataattccatctacattgatggtagccctat  
agaggaaatgggcggtaaaaagggtctaccagtcaaacggtaaatgtcaatgatagttggaatatctggccccaggggac  
atgactacatctatgacaccatttccggagatccaatgagtcacgttttttttactgggtgcatactttatgtctaccaagtgc  
agctgcattgacccaaaaagatagatgtggacgaagatgaaaacatgaatgggatcgaaccgtacattggttggctgtacc  
5 gatgtgtactcaaagacatcaacgttgaagcctgtagggttccttccacctgaagtgtccatcagccttagcgacgtcgc  
atgttccggtataccgttagacattagtagaaccatgaaagagaaatctagcgcaaccgattcaggcccaaaggcataga  
gtcatgcttgaatacaagaaatgtccttaaacatgttgaagttgtttcttctggcagcacgtgtgcgaagtttctgcttagtt  
ctatgatgttcaatgttcaggaaatgttctgttacaataagtttcagcgcatttccagtaaaccttcaattgaaaagggtcag  
caaagatttctacggatacatcaccgttgcgaagatgaatgtgtctgcagtggagttgaaagtgacaaaaatgaagatattt  
10 cgaccagaatgagttcacagttgttttcttaaggaaagtggaattgtggaatactggctctgttcgctcctcttgaatttctt  
gtcgaatgcgtacttccaggtctcattgaaccgttgaagagatcaacaaaccgtgctgttggttggcattctcttcttcat  
ttatagtgaaccttactgagtgtcctgtaaaaagagagagaatgggaactgaacggctatatcatcgcctccaatctt  
ggtatagtttaattatggtagcatgaagcacattctctaaaggatcgtgtgcaatagactcggcgtccactttgtaatagcagt  
gatgagatttggcaatatttctgcataaacctgttctatggcaaaaccagattgcgcacacttcttaatagatagtcggta  
15 aacgcatgcgaaaaagcggtaagaagaccaattggcatacgagccatttcaaaaggaccatctcgaggtaccgatccga  
gacggccggctgggccacgtgaattcgactgctattatctctgtgtgtatgtgtattgggcaacaagagcttgtgaatctgt  
ccttataaaagacaccggaagaggcaagattgggtagtagactaattagtgaggggcaactggttaggggttttgctggc  
ttattatagtgtcagcgatactatacaatctacgatgcccgatgtcggcatacagcacattattagatccaaatttcttaattc  
ctattcacgttatatatattaaccagaatttatccgttgaattgcttaatcgacttctccacgggttggtccgctgtctccgctggg  
20 gcagcaccacctggggcaccgcccagctgcggcagctccgccagggaacctccagggaagccaccaggagctccacc  
aggagctccaccagcagctccatagaactgtctcattattgggttagcaacttctccaattcctttgcctgtccttgactcgtc  
tgtggaagcagattgtgactcatcatagatctgatctcat

**SEQ ID NO: 24 pGAPZAgIsIIHDEL**

25 tcgagatggctcttttgaatggctcgtatgccaatggctcttaccgcttttcgcatgcgttaccgactatctattaaagaa  
gtgtgcgcaatctgggtttgccatagaacagggttatgcagaaaatattgccaatctcatcactgctattacaaagtgga  
cgccgagcttattgcacacgatcctttagagaatgtgcttcatgctaccataattaaaactataccaagattggagggcgatga  
tatagccgttcagttccattctctctctttttacaggatcactcagtaagggtcactataaatgagaaagagagaatgccaa  
ccaacagcagcgggttgatctcttcacaacgggtcaatgagacctggaagtacgcattcgacaagaaattcaagagga

ggcgaacaggaccagtattccacaattccacttccttaagcaaaaacaaactgtgaactcattctggcgaaaatatcttcatt  
ttgtcactttcaaactccactgcagacacatttcattctcgaaacgggtgatgtatccgtagaaatcttctgtaacctttcaattg  
aaagttactggcaaaatgcgctgaaactattgtaaacgagcaaaatttcctgaacattgaacatcatagaactaagcagga  
aaacttcgcacacgtgctgccagaagaacaactttcaacatgtttaaggacaatttctgtattcaaagcatgactctatgcctt  
5 tggggcctgaatcggttgcgctagatttctcttcattgggttctactaatgtctacgggtataccggaacatgcgacgtcgctaag  
gctgatggacacttcaggtggaaaggaaccctacaggctttcaacgttgatgtctttagtacaacatcgggtaccagccaac  
caatgtacgggtcgatccattcatgtttcatcttcgtccacatctatctttgggtcaatgcagctgacacttgggtagacataa  
agtatgacaccagtaaaaaataaaacgatgactcattggatctccgaaaatgggtgcatagatgtatgatgtccctggggcca  
gatattccaactatcattgacaaattaccgatttgactggtagacccttttaccgcccatttctctatagggtaccatcaatgta  
10 gatggaattataatgatgagatggacgttctcagatggactctcagatggatgctcatatgattccttacgattttatttgggtgg  
acttggagtatacgaacgacaaaaaatattttacttggaaagcagcactccttcccaatccaaaaaggctgttatccaaattaaa  
aaagttgggtagaaatcttctgtactaatcgatctcatttaaagaagattatgaaatcagtacagggttaattaatgaaat  
gtagcagtcaaggatcacaatggaaatgactatgtaggctattgctggccaggttaattctatattgattgataccataagcaaa  
tatggccaaaagatttggaaagtccttttgaacgggttatggatctgccggctgatttaactaatttattcatttggatgatga  
15 acgagccttcgatttctgatggcccagagaccacagctccaaaagatttgattcacgacaattacattgaggaaagatccgtc  
cataacatatatggcttatcagtgcataagctacttacgacgcaataaaatcgatttattcaccatccgataagcgctcttct  
tctaacaagggtctttttgccggctctcaacgtactgctgccacatggactgggtgacaatgtggccaattgggattacttaaag  
atttccattcctatggttctgtcaacaacattgctggatgccatttataggagccgacatagctggcttctgaggatcctac  
acctgaattgattgcacgttgggtaccaagcgggcttatgggtaccatttttagagcacacgcccatatagacaccaagagaa  
20 gagaaccatacttattcaatgaacctttgaagtcgatagtacgtgatattatccaattgagatatttctgctacctaccttataca  
ccatgttccataaatcaagtgtcactggatttccgataatgaatccaatgtttattgaacaccctgaatttgcgaattgtatcatat  
cgataaccaattttactggagtaattcaggtctattagtcaaacctgtcacggagcctgggtcaatcagaacggaaatggtttc  
ccaccgggtatattctatgaattcgcattttacactctttataaacaatgggtactgattgatagaaaagaatatttctgcacat  
tggataaaaattccattatttgaaggcgggtcacattatcactatgaagataagtatagaagatcttcaatgttaataaaaaac  
25 gatccatatgtaatagttatagcccctgataccgaggggacgagccgttgagatctttatgttgatgatggagaacttttggct  
accaaagagggtgagtacgtagaactcagttcatttctgaaaacaataaccttaaaaaatgttcgaagtcatttcccagaattt  
gacaggcattcaccacaatacttggaggaataccaatattgaaaaatcattatcgcaagaataattacaacacaacataac  
gttgaaagacagtattaaagtaaaaaaatggcgaagaaggttcattgccgactagatcgtcatatgagaatgataataaga  
tcaccattctaacctatcgcttgacataactgaagattgggaagttattttggggccgaacaaaactcatctcagaagagg



atctgaatagcggcgtcgaccacgacgaactgtgagtttagccttagacatgactgttcctcagttcaagttgggcacttacg  
 agaagaccgggtcttgctagattctaataagaggatgtcagaatgccatttgcctgagagatgcaggcttcattttgatacttt  
 ttattgtaacctatatagtataggattttttgtcattttgttcttctcgtacgagcttgctcctgatcagcctatctcgagctgat  
 gaatacttctggttaggggtttgggaaaatcattcagtttgatgttttcttggtatttccactcctcttcagagtacagaagatt  
 5 aagtgagaccttcgttctgcggatccccacacaccatagcttcaaatgtttctactcctttttactcttcagattttctcgga  
 ctccgcgcacgcccgtaccacttcaaacacccaagcacagcatactaaattttccctcttcttctctaggggtgcgttaatta  
 cccgtactaaagggttggaaaagaaaaagagaccgctcgttcttttctcgtcgaaaaaggcaataaaaaattttatcacg  
 ttctttttcttgaaatttttttttagtttttcttcttcagtgcacctcatgatatttaagtaataaacggtcttcaattctcaagttc  
 agtttcattttctgttctattacaacttttttacttctgttcatttagaaagaaagcatagcaatctaataaggcggtgttgac  
 10 aattaatcatcgccatagatatcggcatagataatacgcagaaggtgaggaactaaacctggccaagtgaccagtgccgt  
 tccgggtgctcaccgcgcgcgacgtcggcgagcgggtcagttctggaccgaccggtcgggttctccgggacttcgtgg  
 aggacgacttcgccggtgtggtccgggacgacgtgacctgttcatcagcgcggtccaggaccaggtggtgccggacaa  
 cacctggcctgggtgtgggtgcgcggcctggacgagctgtacgccgagtggtcggaggctgtgtccacgaactccgg  
 gacgcctccgggcccggccatgaccgagatcggcgagcagccgtggggcgagggttcgcctgcgcgacccggccg  
 15 gcaactgcgtgcacttcgtggccgaggagcaggactgacacgtccgacggcgcccacgggtccaggcctcggagat  
 ccgtcccccttttcttctgcatatcatgtaattagttatgtcacgcttacattcacgccctccccccacatccgctctaaccgaa  
 aaggaaggagttagacaacctgaagtctaggtccctattttttatagttatgttagtattaagaacgttatttatatttcaaatt  
 ttctttttttctgtacagacgcgtgtacgcatgtaacattatactgaaaaccttgccttgagaaggtttgggacgctcgaaggct  
 ttaatttgcaagctggagaccaacatgtgagcaaaaggccagcaaaaggccagggaaccgtaaaaaggccgcgttgctgg  
 20 cgttttccataggtcggccccctgacgagcatcacaaaaatcgacgctcaagtcagaggtggcgaaacccgacagga  
 ctataaagataccaggcgtttccccctggaagctccctcgtgcgctcctgttccgacctcggcgttaccggatacctgtcc  
 gcctttctcccttcgggaagcgtggcgctttctcaatgtcacgctgtaggtatctcagttcgggtgtaggtcgttcgctccaagc  
 tgggctgtgtgcacgaacccccgttcagcccagccgctgcgccttatccggttaactatcgttcttgagccaacccggttaag  
 acacgacttatgccactggcagcagccactggtaacaggattagcagagcaggtatgtaggcgggtgtacagagttctt  
 25 gaagtgggtggcctaactacggctacactagaaggacagtatttgggtatctgcgctctgtgaagccagttaccttcggaaaaa  
 gagttggtagctcttgatccggcaaacaaaccaccgctggtagcggtgtttttgttgcaagcagcagattacgcgcaga  
 aaaaaaggatctcaagaagatcctttgatctttctacggggtctgacgctcagtggaacgaaaactcacgttaagggttttg  
 gtcagtcagatgagatcagatctttttgtagaatgtcttggtgtcctcgtccaatcaggtagccatctctgaaatatctggctccgt  
 tgcaactccgaacgacctgctggcaacgtaaaattctccgggtgtaaaactaaatgtggagtaatggaaccagaaacgtctc

ttcccttctctctctccaccgcccgttaccgtccctaggaattttactctgctggagagcttctctacggcccccttgccagc  
aatgctcttcccagcattacgttgccggtaaaacggaggtcgtgtacccgacctagcagcccagggtggaaaagtccc  
gccgtcgtggcaataatagcgggcggacgcatgcatgagattattgaaaccaccagaatcgaatataaaaggcgaac  
accttcccaattttggttctcctgacccaaagactttaatttaattttgtccctatttcaatcaattgaacaactatttcgaaa  
5 cgaggaattcacgtggcccagccggccgtctcggatcgggtacc

**SEQ ID NO: 25** pGAPADE1glsIIHDEL

cgctactcacctctttggtagccaaaagtcttccatcatcaacataaagatctccaacggctcgtccctcgggtatcaggggctat  
aactattacatatggatcgttttcattaacattgaagatcttctatacttattcttcatagtgataatgtgaccgccttcaataataa  
10 tggaaattta'ccaatggtgcagaaatattctttctatcaaatcagtaccattgtttataaaagagtgtaaagatgcgaattcatag  
aatataccgggtgggaaaaccatttccgtttctgattgaccaggctccgtgacagggttgactaataagacctgaattactccag  
taaaattggtatcgtatgatacaattcagcaaatcagggtgttcaataaacattggattcattatcggaatccagtgcact  
tgatttatgaaacatggtgtataaggtaggtagcaggaaatctcaattggataatatcacgtactatcgacttcaaagggtcat  
tgaataagtatggttctctctctgtgtctatattggcggtgtgtcttaaaaaatgggtaccataagcccgttgggtaccaacgt  
15 gcaatcaattcaggtgtaggatcctcagcaaaaggcagctatgtcggctcctataaatggcataccagcaatgtgtttgacag  
aaccataggaatggaaatctttaagtaatcccaattggccacattgtcaccagtccatgtggcagcagtacgttgagagccg  
gcaaaaaaagcccttggtagaaggaaaggacgcttaccgatggtgaataaatcgattttattgcgtcgtgaagtagcttcatgc  
actgatagaccatatattgtatggacggatcttctcctcaatgtaattgtcgtgaatcaaatcttttgagctgtgtgtctctgggcca  
tcgaaaatcgaggctcgttcatatcattccaaatgaataaattagttaaatcagccggcagatccataaacggttcgaaaaag  
20 gacttccaaatcttttgccatatttgcttatggtatcaatccatatagaattacctggccagcaatgacctacatagtcatttccat  
tgtgatccttgactgctacatttccatttaattaccctgtcactgatttcataatcttctttaaattgaggatcgattagtacgacaaga  
tttctacccaacttttttaatttgataacagcccttttgattgggaaaggagtgtgtcttccaagtaaaatatttttgcgttcgta  
tactccaagtcacaaataaaatcgtgaaggaaatcatatgagcatccatctgagagtccactgtgagaacgtccatctcatc  
attataattccatctacattgatgtaccctatagaggaaatgggcggtaaaaagggtctaccagtc aaatcggtaaatgtca  
25 atgatagttggaatatctggccccaggacatgactacatctatgacaccatttccggagatccaatgagtcacgttttattttta  
ctggtgtcactttatgtctacccaagtgtcagctgcattgacccaaaagatagatgtggacgaagatgaaaacatgaatgg  
gatcgaaccgtacattggttggctgtgtaccgatgtgtactcaagacatcaacgttgaaggcctgtagggttcccttccacc  
tgaagtgtccatcagcccttagcgacgtcgcattgtccggtataccgtagacattagtagaaccatgaaagagaaatctagc  
gcaaccgattcaggcccaaggcatagatgctttgaatacaagaaattgtccttaaacatgttgaaagtgtttcttctg

gcagcacgtgtgcgaagttttcctgcttagttctatgatgttcaatgttcaggaaatttgctcgtttacaataagtttcagcgcat  
ttgccagtaaactttcaattgaaaagggttcagcaaagatttctacggatacatcaccgtttcgaagatgaaatgtgtcgcagt  
gagtttgaaagtacaaaaatgaagatatttcgaccagaatgagttcacagttgttttgcttaagggaagtggaaatgtggaat  
actggctcgttcgcctcctcttgaaatttctgtcgaatgcgtacttccaggtctcatgaaccgttggaagagatcaacaac  
5 cgctgctgttggtggcattctcttctcatttatagtgaaacctactgagtgatcctgtaaaaaagagagagagaatgggaa  
ctgaacggctatatcatcgccctccaatctggtagtttaattatggtagcatgaagcacattctctaaggatcgtgtgcaa  
tagactcggcgtccacttgaatagcagtgatgagatttggaatatttctgcataaacctgtttctatggcaaaaccagat  
tgcgcacacttcttaatagatagtcggtaaacgcattgcgaaaaagcggtaaagaagaccaattggcatacagccatttca  
aaaggaccatctcgaggtaccgatccgagacggccggctggggccacgtgaattcctcgtttcgaaatagtgttcaattgatt  
10 gaaatagggacaaataaattaaatftaaagtctttgggtcaggagaaacaaaattgggaaaggtgttcgcttttatattcgat  
tctggtggttccaataatctcatgacatgcgtccgccgctattattgccagcgacggccgggacttttccatccctgggctg  
ctaggtcgggtacacgacctccgtttaccgcgaacgtaatgctgggaagagcattgctgcaagggggccgtagaagaag  
ctctccagcagagtaaaatttctagggacggtaacggcggtggaaggagagagaagggaagagacgtttctggttcca  
ttactccacatttaagttttaccggagaattttacgttgccagcaggtcgttcggagtgcacaggagccagatatttcagag  
15 atggtctacctgattggacgaggacaccaagacatttctacaaaaagatctgatctcatcgaccggctgcattaatgaatcgg  
ccaacgcgcggggagaggcggtttgcgtattgggcgctcttccgcttctcgtcactgactcgtcgctcggctcgttcgg  
ctgcggcgagcgggtatcagctcactcaaaaggcggtataacaggttatccacagaatcaggggataacgcaggaaagaacat  
gtgagcaaaaggccagcaaaaggccaggaaccgtaaaaaggccggttgcgtggcgttttccataggctccgccccctg  
acgagcatcacaaaaatcgacgtcaagtcagaggtggcgaaacccgacaggactataaagataaccaggcgtttccccct  
20 ggaagctccctcgtgcgtctcctgttccgacctgccgcttaccggatacctgtccgccttctccctcgggaagcgtggc  
gctttctcatagctcacgctgtaggtatctcagttcgggtgtaggtcgttcgtccaagctgggctgtgtgcagcaacccccgt  
tcagcccagccgctgcgccttatccggtaactatcgtcttagtccaacccggtgaagacacgacttatcgccactggcagca  
gccactggtaacaggattagcagagcgaggtatgtaggcggtgtacagagttcttgaagtgtggcctaactacggctac  
actagaaggacagtatttggtagtctgcgtctgctgaagccagttaccttcggaaaaagagttggtagctcttgatccggcaa  
25 acaaaccaccgctggtagcgggtgtttttgttgcaagcagcagattacgcgcagaaaaaaggatctcaagaagatcctt  
tgatcttttctacgggggtctgacgctcagtggaacgaaaactcacgttaagggttttggatgagattatcaaaaaggatctt  
cacctagatccttttaattaaaaatgaagttttaaataatctaaagtatatatgagtaaaacttggtctgacagttaccaatgctta  
atcagtgaggcacctatctcagcgatctgtctatttcgttcatccatagttgcctgactccccgtcgtgtagataactacgatac  
gggaggggttaccatctggccccagtgctgcaatgataccgcgagacccacgctcaccggctccagatttatagcaataa

accagccagccggaaggggccgagcgcagaagtggcctgcaactttatccgcctccatccagctctattaattgttgcggg  
aagctagagtaagtagttcggcagtaatagtttgcgaacgttggtgccattgctacaggcatcgtgggtgcacgctcgtcgtt  
tggatggcttcattcagctccggttcccaacgatcaaggcgagttacatgatcccccattgttgcaaaaaagcggttagctc  
cttcggctcctccgacgttgtcagaagtaagttggccgagtggtatcactcatggttatggcagcactgcataattctcttactg  
5 tcatgccatccgtaagatgcttttctgtgactggtagtacicaaccaagtcattctgagaatagtgatgcggcgaccgagtt  
gctcttggccggcgtcaatacgggataataccgcccacatagcagaactttaaaagtgcctcatcattggaaaacgttctcg  
ggcgcaaaactctcaaggatcttaccgctgttgagatccagttcgtatgaaccactcgtgcaccaactgatcttcagcatct  
ttactttcaccagcggttctgggtgagcaaaaacaggaaggcaaaatgccgcaaaaaaggggaataaggcgacacgggaa  
atgttgaatactcatactcttcttttcaatagctccaaggcaacaaattgactactcagaccgacattcgttatigattttaa  
10 atcaacgataaacggatggttacttgaatgatttcactttatgatcattgtttactaattacctaataaggattttatatggaattgg  
aagaataagggaatttcagatgcttgaaaaaggcgaggagggtactaatcattcaagcccatttcttgcagtaattgcttca  
taagcttcaatatacttttcttactcttgatagcaatttctgcacccatggctacgccctcttggccattcaatccgttggccgtaaa  
ccaatctctgagaaactgcttatcgtaaactcttgcgatttaccacttggtaagtcttttgattccaaaatctagaagaatctgg  
agttaaaactctactagtaccaattcattgtttcgtccagtccaaaatcgaatttcgtatcagcaataatgatcccttcaaa  
15 agggcgaaagtttttgcagcagaataacaactcgaccgccttgacagcgaccttctcacaatgtctttacctacaatctcagca  
gcttgttcaatagagatgtttcatcgtgttcaccctgttcagcttctgttgaaaggttgaaaaatcgaggttgaaaggcgctcgt  
ctcttgaagggttctcgtttcaaccttgactccatggacagttttgagttctgtactcttccatgcacttccagtgatgtaacctc  
tgacaatggcttccaaaggatcagctgtgtttttactatcaaggatcgtccctctaattgagatttgtatttttctcagacagt  
tttgatggtagtaaaagcaagacttccctgtcattagaagcaaccaaattgattctttatgtagggtgccaaaaatcaaacag  
20 aaaactgagagctgagtcaaaatcttcccttatcaggaataaccgtttgcataatcacatcgtaaaggagatacggtcagtt  
gcgacgaacagcaagttgttctcatcgactgcataaatgtctctaaccttcccttggcgattaaaggtaggattccgtccagat  
cagtggttcacaatggacatacttgaaggatacagcaaaagtgtgttgaagcgatgacacatggaaagggaatttttcagattt  
cctagagtagtatattggggcggtgaaagttcagatgtttaatgcttaatactcttatacttctcaaacgcccgaagtgttctgc  
caacctgactttttctgaataatgaatcgttcaagtgagattttaaacatgattaagttacgtgatttggcactggataaggtc  
25 gaaaaatatccgtattcataaacgattattggtaaaagtacaaaataccactaattacggagaagccttagtaacagttatcatct  
cttggctgattaacgcttacaatttccattcgccattcagggtgcgcaactgttgggaaggcgatcggtgcgggcctcttcg  
ctattacgccagggcctcgaggcacaacgaacgtctcacttaactctgtactctgaagaggagtgaggaaataccaagaa  
aaacatcaaaactcgaatgattttcccaaacccctaccacaagatattcatcagctgcgagataggctgatcaggagcaagct  
cgtacgagaagaacaaaaatgacaaaaaaatcctatactatataggttacaaataaaaaagtatcaaaaatgaagcctgca

tctctcaggcaaatggcattctgacatcctcttgattagaatctagcaagaccgggtctctcgtaatgcccaactgaactgag  
gaacagtcattgtctaaggctaaaactcacagttcgtcgtggtcgacggcgctattcagatcctctctgagatgagttttgttc  
gggccccaaaaataactcccaatcttcagttatgtcaagcgataggtaagaatggtgatcttattatcattctcatatgacgatc  
tagtcggcaatgaactttctcgccattttttgactttaatactgtctttcaacgttatgttggttgtaaattattcttgcgataatg  
5 atttttcaatattggtattcctcaaagtattgtggtgaatgcctgtcaaattctcgggaatatgacttcgaacatttttaaggattg  
ttttcgaaaatgaactgagtttcta

**SEQ ID NO: 26**

GACGAGATCTTTTTTTCAGACCATATGACCGG  
10

**SEQ ID NO: 27**

GCGGAATTCTTTTCTCAGTTGATTTGTTTGT

**SEQ ID NO: 28**

15 GCG GGT CGA CCA CGA CGA ACT GTG AGT TTT AGC CTT AGA CAT GAC

**SEQ ID NO: 29**

CAG GAG CAAA GCT CGT ACG AG  
20

**We claim:**

1. A vector capable of expressing an  $\alpha$ -1,2-mannosidase or a functional part thereof in a methylotrophic yeast strain, comprising a nucleotide  
5 sequence coding for said  $\alpha$ -1,2-mannosidase or said functional part.
2. The vector of claim 1, wherein said  $\alpha$ -1,2-mannosidase is a protein from a fungal species.
- 10 3. The vector of claim 2, wherein said fungus is *Trichoderma reesei*.
4. The vector of claim 1, wherein said  $\alpha$ -1,2-mannosidase is a protein from a mammalian species.
- 15 5. The vector of claim 4, wherein said  $\alpha$ -1,2-mannosidase is murine  $\alpha$ -1,2-mannosidase IA or IB.
6. The vector of claim 1, wherein said  $\alpha$ -1,2-mannosidase or said functional part is tagged with an ER-retention signal.  
20
7. The vector of claim 6, wherein said ER-retention signal comprises peptide HDEL.
8. The vector of claim 1, wherein the nucleotide sequence coding for  
25 said  $\alpha$ -1,2-mannosidase or said functional part is operably linked to a promoter and a 3' termination sequence.

9. The vector of claim 8, wherein said promoter is the promoter of a gene selected from the group consisting of AOXI, AOXII, GAP, and FLD.

10. A vector selected from the group consisting of  
5 pGAPZMFManHDEL, pGAPZMFManMycHDEL, pPICZBMFManMycHDEL, pGAPZmManHDEL, pGAPZmMycManHDEL, pPIC9mMycManHDEL and pGAPZmMycManHDEL.

11. A vector capable of expressing a glucosidase II or a functional part  
10 thereof in a methylotrophic yeast strain, comprising a nucleotide sequence coding for said glucosidase II or said functional part.

12. The vector of claim 11, wherein said glucosidase II is a protein  
15 from a fungal species.

13. The vector of claim 12, wherein said fungus is *Saccharomyces cerevisiae*.

14. The vector of claim 11, wherein said glucosidase II is a protein  
20 from a mammalian species.

15. The vector of claim 11, wherein said glucosidase II or said functional part is tagged with an ER-retention signal.

16. The vector of claim 15, wherein said ER-retention signal comprises  
25 peptide HDEL.

17. The vector of claim 11, wherein the nucleotide sequence coding for said  $\alpha$ -1,2-mannosidase or said functional part is operably linked to a promoter and a 3' termination sequence.
- 5 18. The vector of claim 17, wherein said promoter is the promoter of a gene selected from the group consisting of AOXI, AOXII, GAP, and FLD.
19. A vector having the designation pGAPZAGLSII, pPICZAGLSII, pAOX2ZAGLSII, pYPTIZAGLSII, pGAPADEglsII, pPICADEglsII,  
10 pAOX2ADEglsII, pYPTIADEglsII, pGAPZAglslIIHDEL and pGAPADEglsIIHDEL.
20. A vector for disrupting the Och1 gene in a methylotrophic yeast strain, comprising a portion of the Och1 gene and a selectable marker gene, wherein said portion of the Och1 gene and said selectable marker gene are linked in such a way  
15 to effect the disruption of the genomic Och1 gene in said methylotrophic yeast strain.
21. A vector having the designation pBLURA5'PpOCH1.
22. A method of reducing the glycosylation on proteins produced from  
20 a methylotrophic yeast, comprising transforming said yeast with any one of the vectors of claims 1-21.
23. The method of claim 22, wherein said yeast is *Pichia pastoris*.
- 25 24. The method of claim 23, wherein said yeast is a *Pichia pastoris* strain selected from GS115 (NRRL Y-15851), GS190 (NRRL Y-18014), PPF1 (NRRL Y-18017), PPY12-OH, yGC4, or derivatives thereof.



25. The method of claim 22, 23 or 24, wherein said yeast has been genetically engineered to express a heterologous protein.

26. A genetically engineered strain of a methylotrophic yeast, wherein  
5 said strain is transformed with at least one of the vectors of claims 1-21.

27. A method of reducing the glycosylation of a heterologous glycoprotein expressed from a methylotrophic yeast, comprising transforming cells of said methylotrophic yeast with at least one of the vectors of claims 1-21, and  
10 producing said glycoprotein from the transformed cells.

28. A method of producing a glycoprotein with reduced glycosylation in a methylotrophic yeast, comprising transforming cells of said methylotrophic yeast with at least one of the vectors of claims 1-21 and with a nucleotide sequence capable of expressing said glycoprotein in said yeast, and producing said glycoprotein from the  
15 transformed cells.

29. A glycoprotein produced by the method of claim 27 or 28.

20 30. The glycoprotein of claim 29, wherein said glycoprotein has a reduced immunogenicity as relative to the glycoprotein produced from a wild type strain of said methylotrophic yeast.

31. The glycoprotein of claim 29, wherein said glycoprotein is suitable  
25 for use in human therapeutics.

32. A kit comprising any of the vectors of claims 1-21.

33. The kit of claim 32, further comprising a methylotrophic yeast strain.

34. A kit comprising the methylotrophic yeast strain of claim 26.

5




FIGURE 1

L929 (cycloheximide+  $Ti^{-}$  induced) mouse cDNA library,  
Group 6, average insert length : 2000 bp.




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6






FIGURE 7




FIGURE 8

### Pichia OCH1 Genomic fragment




FIGURE 9




FIGURE 10




FIGURE 11

## GlsII Pichia expression vector pGAPZ(A)glsII




FIGURE 12


GlsII Pichia expression vector pAOX2Z(A)gls<sup>II</sup>

FIGURE 13

## GlsII Pichia expression vector pPICZ(A)glsII




FIGURE 14



## GlsII Pichia expression vector pYPT1Z(A)glsII




FIGURE 15

## GlsII Pichia expression vector pGAPADE1glsII




FIGURE 16

## GlsII Pichia expression vector pAOX2ADE1glsII




FIGURE 17

## GlsII Pichia expression vector pPICADE1glsII




FIGURE 18

## GlsII Pichia expression vector pYPT1ADE1glsII




FIGURE 19




FIGURE 20

## GlsII Pichia expression vector pGAPADE1glsIIHDEL




FIGURE 21

# Glucosidase II assay on commercially available alpha-glucosidase




FIGURE 22



## Glucosidase II assay on heterologously expressed Pichia protein




FIGURE 23